arXiv:1601.08136v2 [math.PR] 1 Aug 2016

FRACTIONAL POISSON FIELDS AND MARTINGALES

By Giacomo Aletti* Nikolai Leonenko† and Ely Merzbach‡

*ADAMSS Center & Università degli Studi di Milano, Italy
†Cardiff University, United Kingdom
‡Bar-Ilan University, Israel

We present new properties for the Fractional Poisson process and the Fractional Poisson field on the plane. A martingale characterization for Fractional Poisson processes is given. We extend this result to Fractional Poisson fields, obtaining some other characterizations. The fractional differential equations are studied. We consider a more general Mixed-Fractional Poisson process and show that this process is the stochastic solution of a system of fractional differential-difference equations. Finally, we give some simulations of the Fractional Poisson field on the plane.

1. Introduction. There are several different approaches to the fundamental concept of Fractional Poisson process (FPP) on the real line. The "Renewal" approach consists of considering the characterization of the Poisson process as a sum of independent non-negative random variables, and instead of the assumption that these random variables have an exponential distribution, we assume that they have the Mittag-Leffler distribution (see [31, 32, 41]). In [6], the renewal approach to the Fractional Poisson process is developed and it is proved that its one-dimensional distributions coincide with the solution to fractionalized state probabilities. Another equivalent approach is using "inverse subordinator", a kind of Fractional Poisson process can be constructed [33].

In [26], following this method, the FPP is generalized and defined, obtaining a Fractional Poisson random field (FPRF) parametrized by points of the Euclidean space $\mathbb{R}^2_+$, as it has been done for Fractional Brownian fields, see, e.g., [18, 23, 29, 21].

The starting point of our extension will be the set-indexed Poisson process which is a well-known concept, see, e.g., [43, 36, 37, 18, 23].

In this paper, we first present a martingale characterization of the Fractional Poisson process. We extend this characterization to FPRF using the concept of increasing path and strong martingales. This characterization permits us to give a definition of a set-indexed Fractional Poisson process. We study the fractional differential equation for FPRF. Finally, we study Mixed-Fractional Poisson processes.

The paper is organized as follows. In the next section, we collect some known results from the theory of subordinators and inverse subordinators, see [9, 45, 46, 35] among others. In Section 3, we prove a martingale characterization of the FPP, which is a generalization of the Watanabe Theorem. In Section 4, another generalization called "Mixed-Fractional Poisson process" is introduced and some distributional properties are studied as well as Watanabe characterization is given. Section 5 is devoted to FPRF. We begin by computing covariance for this process, then we give some characterizations using increasing paths and intensities. We present a Gergely-Yeshow characterization and discuss random time changes. Fractional differential equations are discussed on Section 6.

Finally, we present some simulations for the FPRF.

Other different generalizations of FPP and fields can be found in [47, 48, 8, 38, 39].

2. Inverse Subordinators. This section collects some known resuts from the theory of subordinators and inverse subordinators [9, 45, 46, 35].

2.1. *Subordinators and their inverse.* Consider a non-decreasing Lévy process $L = \{L(t),\ t \geq 0\}$, starting from 0, which is continuous from the right with left limits (cadlag), continuous in probability, with independent and stationary increments. Such a process is known as a Lévy subordinator with Laplace exponent

$$\phi(s) = \mu s + \int_{(0,\infty)} (1 - e^{-sx})\Pi(dx), \quad s \geq 0,$$

where $\mu \geq 0$ is the drift and the Lévy measure Π on $\mathbb{R}_+ \cup \{0\}$ satisfies

$$\int_0^\infty \min(1, x)\Pi(dx) < \infty.$$

This means that

$$\mathrm{E}e^{-sL(t)} = e^{-t\phi(s)},\ s \geq 0.$$

Consider the inverse subordinator $Y(t),\ t \geq 0$, which is given by the first-passage time of L :

$$Y(t) = \inf\{u \geq 0 : L(u) > t\}, t \geq 0.$$

The process $Y(t),\ t \geq 0$, is non-decreasing and its sample paths are a.s. continuous if and its sample paths are a.s. right continuous if L is strictly increasing.

We have

$$\{(u_i, t_i) \colon L(u_i) < t_i, i = 1, \ldots, n\} = \{(u_i, t_i) \colon Y(t_i) > u_i, i = 1, \ldots, n\},$$

and for any $z > 0$

$$\mathrm{P}\{Y(t) > x\} = \mathrm{P}\{L(x) \leq t\} = \mathrm{P}\left\{e^{-zL(x)} \geq e^{-zt}\right\} \leq \exp\{tz - x\phi(z)\},$$

which implies that for any $p > 0, \mathrm{E}Y^p(t) < \infty$, since

$$\mathrm{E}Y^p(t) = p\int_0^\infty x^{p-1}(1 - \mathrm{P}\{Y(t) \leq x\})dx \leq pe^{tz}\int_0^\infty x^{p-1}e^{-x\phi(z)}dx < \infty.$$

Let $U(t) = \mathrm{E}Y(t)$ be the renewal function. Since

$$\tilde{U}(s) = \int_0^\infty U(t)e^{-st}dt = \frac{1}{s\phi(s)},$$

then $\tilde{U}$ characterizes the inverse process Y, since ϕ characterizes L.

We get a covariance formula [45, 46]

$$\mathrm{Cov}(Y(t), Y(s)) = \int_0^{\min(t,s)} (U(t_1 - \tau) + U(t_2 - \tau))dU(\tau) - U(t_1)U(t_2).$$

The most important example is considered in the next section, but there are some other examples.

2.2. *Inverse stable subordinators.* Let $L_\alpha = \{L_\alpha(t), t \geq 0\}$, be an $\alpha-$stable subordinator with $\phi(s) = s^\alpha, 0 < \alpha < 1$. The density of $L_\alpha(1)$ is of the form [44]

$$g_\alpha(x) = \frac{1}{\pi}\sum_{k=1}^{\infty}(-1)^{k+1}\frac{\Gamma(\alpha k+1)}{k!}\frac{1}{x^{\alpha k+1}}\sin(\pi k\alpha) = \frac{1}{x}W_{-\alpha,0}(-x^{-\alpha}). \tag{2.1}$$

Here we use the Wright' s generalized Bessel function (see, e.g., [17])

$$W_{\gamma,\beta}(z) = \sum_{k=0}^{\infty}\frac{z^k}{\Gamma(1+k)\Gamma(\beta+\gamma k)}, \quad z \in \mathbb{C}, \tag{2.2}$$

where $\gamma > -1$, and $\beta \in \mathbb{R}$. The set of jump times of L_α is a.s. dense. The Lévy subordinator is strictly increasing, since the process L_α admits a density.

Then the inverse stable subordinator

$$Y_\alpha(t) = \inf\{u \geq 0 : L_\alpha(u) > t\}$$

has density [35, p.110]

$$f_\alpha(t,x) = \frac{d}{dx}\mathrm{P}\{Y_\alpha(t) \leq x\} = \frac{t}{\alpha}x^{-1-\frac{1}{\alpha}}g_\alpha(tx^{-\frac{1}{\alpha}}), \quad x > 0, \quad t > 0. \tag{2.3}$$

The Laplace transform of the density $f_a(t,x)$ is

$$\int_0^\infty e^{-st}f_a(t,x)dt = s^{\alpha-1}e^{-xs^\alpha}, \quad s \geq 0, \tag{2.4}$$

Its paths are continuous and nondecreasing. For $\alpha = 1/2$, the inverse stable subordinator is the running supremum process of Brownian motion, and for $\alpha \in (0, 1/2]$ this process is the local time at zero of a strictly stable Lévy process of index $\alpha/(1-\alpha)$.

Let

$$E_\alpha(z) = \sum_{k=0}^{\infty}\frac{z^k}{\Gamma(\alpha k+1)}, \ \alpha > 0, \ z \in \mathbb{C} \tag{2.5}$$

be the Mittag-Leffler function [17], and recall the following:

i) The Laplace transform of function $E_\alpha(-t^\alpha)$ is of the form

$$\int_0^\infty e^{-st}E_\alpha(-\lambda t^\alpha)dt = \frac{s^{\alpha-1}}{\lambda+s^\alpha}, \quad 0 < \alpha < 1, \ t \geq 0, \Re(s) > |\lambda|^{1/\alpha}.$$

(ii) The function $E_\alpha(\lambda t^\alpha)$ is an eigenfunction at the the fractional Caputo-Djrbashian derivative D_t^α with eigenvalue λ [35, p.36]

$$\mathrm{D}_t^\alpha E_\alpha(\lambda t^\alpha) = \lambda E_\alpha(\lambda t^\alpha), \quad 0 < \alpha < 1, \lambda \in \mathbb{R},$$

where D_t^α is defined as (see [35])

$$\mathrm{D}_t^\alpha u(t) = \frac{1}{\Gamma(1-\alpha)}\int_0^t \frac{du(\tau)}{d\tau}\frac{d\tau}{(t-\tau)^\alpha}, \quad 0 < \alpha < 1. \tag{2.6}$$

Proposition 2.1 (See [9, 45, 46]). *The α-stable inverse subordinators satisfy the following properties:*

(i)

$$\mathrm{E}e^{-sY_\alpha(t)} = \sum_{n=0}^{\infty} \frac{(-st^\alpha)^n}{\Gamma(\alpha n+1)} = E_\alpha(-st^\alpha), \quad s>0.$$

(ii) Both processes $L_\alpha(t), t\geq 0$ and $Y_\alpha(t)$ are self-similar

$$\frac{L_\alpha(at)}{a^{1/\alpha}} \stackrel{d}{=} L_\alpha(t), \ \frac{Y_\alpha(at)}{a^\alpha} \stackrel{d}{=} Y_\alpha(t), \quad a>0.$$

(iii) For $0<t_1<\cdots<t_k$,

$$\frac{\partial^k \mathrm{E}(Y_\alpha(t_1)\cdots Y_\alpha(t_k))}{\partial t_1\cdots\partial t_k} = \frac{1}{\Gamma^k(\alpha)} \frac{1}{[t_1(t_2-t_1)\cdots(t_k-t_{k-1})]^{1-\alpha}}.$$

In particular,
(A)

$$\mathrm{E}Y_\alpha(t) = \frac{t^\alpha}{\Gamma(1+\alpha)}; \mathrm{E}[Y_\alpha(t)]^\nu = \frac{\Gamma(\nu+1)}{\Gamma(\alpha\nu+1)} t^{\alpha\nu}, \quad \nu>0;$$

(B)

$$\mathrm{Cov}(Y_\alpha(t), Y_\alpha(s)) = \frac{1}{\Gamma(1+\alpha)\Gamma(\alpha)} \int_0^{\min(t,s)} \left((t-\tau)^\alpha + (s-\tau)^\alpha\right) \tau^{\alpha-1} d\tau - \frac{(st)^\alpha}{\Gamma^2(1+\alpha)}. \quad (2.7)$$

2.3. *Mixture of inverse subordinators.* This subsection collects some results from the theory of inverse subordinators, see [45, 46, 35, 5, 27].

Different kinds of inverse subordinators can be considered.

Let L_{α_1} and L_{α_2} be two independent stable subordinators. The mixture of them $L_{\alpha_1,\alpha_2} = \{L_{\alpha_1,\alpha_2}(t), t\geq 0\}$ is defined by its Laplace transform: for $s\geq 0$, $C_1+C_2=1$, $C_1\geq 0$, $C_2\geq 0$, $\alpha_1<\alpha_2$,

$$\mathrm{E}e^{-sL_{\alpha_1,\alpha_2}(t)} = \exp\{-t(C_1 s^{\alpha_1} + C_2 s^{\alpha_2})\}. \quad (2.8)$$

It is clear that

$$L_{\alpha_1,\alpha_2}(t) = (C_1)^{\frac{1}{\alpha_1}} L_{\alpha_1}(t) + (C_2)^{\frac{1}{\alpha_2}} L_{\alpha_2}(t), \quad t\geq 0,$$

is not self-similar, unless $\alpha_1=\alpha_2=\alpha$, since $L_{\alpha_1,\alpha_2}(at) =^d (C_1)^{\frac{1}{\alpha_1}} a^{\frac{1}{\alpha_1}} L_{\alpha_1}(t) + (C_2)^{\frac{1}{\alpha_2}} a^{\frac{1}{\alpha_2}} L_{\alpha_2}(t)$. This expression is equal to $a^{\frac{1}{\alpha}} L_{\alpha_1,\alpha_2}(t)$ if and only if for any $t>0$ $\alpha_1=\alpha_2=\alpha$, in which case the process L_{α_1,α_2} can be reduced to the classical stable subordinator (up to constant).

The inverse subordinator is defined by

$$Y_{\alpha_1,\alpha_2}(t) = \inf\{u\geq 0 : L_{\alpha_1,\alpha_2}(u) > t\}, \quad t\geq 0. \quad (2.9)$$

We assume that $C_2\neq 0$ without loss of generality (the case $C_2=0$ reduces to the previous case of single inverse subordinator).

It was proved in [27] that

$$\tilde{U}(t) = \frac{1}{(C_1 s^{\alpha_1} + C_2 s^{\alpha_2})s}, U(t) = \frac{1}{C_2} t^{\alpha_2} E_{\alpha_2-\alpha_1,\alpha_2+1}(-\tfrac{C_1}{C_2} t^{\alpha_2-\alpha_1}), \quad (2.10)$$

where $E_{\alpha,\beta}(z)$ is the two-parametric Generalized Mittag-Leffler function ([15, 17])

$$E_{\alpha,\beta}(z) = \sum_{k=0}^{\infty} \frac{z^k}{\Gamma(\alpha k+\beta)}, \ \alpha>0, \beta>0, \ z\in\mathbb{C}.$$

Also for the Laplace transform of the density $f_{\alpha_1,\alpha_2}(t,u) = \frac{d}{du}\mathrm{P}\{Y_{\alpha_1,\alpha_2}(t) \leq u\}$, $u \geq 0$, of inverse subordinator $Y_{\alpha_1,\alpha_2} = \{Y_{\alpha_1,\alpha_2}(t), t \geq 0\}$, we have the following expression [34]:

$$\tilde{f}_{\alpha_1,\alpha_2}(s,u) = \int_0^\infty e^{-st} f_{\alpha_1,\alpha_2}(t,u)dt = \frac{1}{s}[C_1 s^{\alpha_1} + C_2 s^{\alpha_2}]e^{-u[C_1 s^{\alpha_1} + C_2 s^{\alpha_2}]}, \quad s \geq 0, \tag{2.11}$$

and the Laplace transform of $\tilde{f}$ is given by

$$\int_0^\infty e^{-pu}\tilde{f}_{\alpha_1,\alpha_2}(s,u)du = \frac{\phi(s)}{s(p+\phi(s))} = \frac{C_1 s^{\alpha_1 - 1} + C_2 s^{\alpha_2 - 1}}{p + C_1 s^{\alpha_1} + C_2 s^{\alpha_2}}, \quad p \geq 0. \tag{2.12}$$

From [5, Theorem 2.3] we have the following expression for $u \geq 0, t > 0$:

$$f_{\alpha_1,\alpha_2}(t,u) = \frac{C_1}{\lambda t^{\alpha_1}} \sum_{r=0}^\infty \frac{1}{r!}(-\frac{C_2\,|u|}{\lambda t^{\alpha_2}})^r W_{-\alpha_1, 1-\alpha_2 r - \alpha_1}(-\frac{C_1\,|u|}{\lambda t^{\alpha_1}}) +$$

$$+ \frac{C_2}{\lambda t^{\alpha_2}} \sum_{r=0}^\infty \frac{1}{r!}(-\frac{C_1\,|u|}{\lambda t^{\alpha_1}})^r W_{-\alpha_2, 1-\alpha_1 r - \alpha_2}(-\frac{C_2\,|u|}{\lambda t^{\alpha_2}}). \tag{2.13}$$

One can also consider the tempered stable inverse subordinator, the inverse subordinator to the Poisson process, the compound Poisson process with positive jumps, the Gamma and the inverse Gaussian Lévy processes. For additional details see [45, 46, 27].

3. Fractional Poisson Processes and Martingales.

3.1. *Preliminaries.* The first definition of FPP $N_\alpha = \{N_\alpha(t), t \geq 0\}$ was given in [31] (see also [32]) as a renewal process with Mittag-Leffler waiting times between the events

$$N_\alpha(t) = \max\{n : T_1 + ... + T_n \leq t\} = \sum_{j=1}^\infty \mathbb{1}_{\{T_1 + ... + T_j \leq t\}}, \qquad t \geq 0,$$

where $\{T_j\}$, $j = 1, 2, ...$ are iid random variables with the strictly monotone Mittag-Leffler distribution function

$$F_\alpha(t) = \mathrm{P}(T_j \leq t) = 1 - E_\alpha(-\lambda t^\alpha), \qquad t \geq 0, 0 < \alpha < 1, \quad j = 1, 2, \ldots$$

The following stochastic representation for FPP is found in [33]:

$$N_\alpha(t) = N(Y_\alpha(t)), \quad t \geq 0, \quad \alpha \in (0,1),$$

where $N = \{N(t), t \geq 0\}$, is the classical homogeneous Poisson process with parameter $\lambda > 0$, which is independent of the inverse stable subordinator Y_α. One can compute the following expression for the one-dimensional distribution of FPP:

$$\begin{aligned}
\mathrm{P}(N_\alpha(t) = k) = p_k^{(\alpha)}(t) &= \int_0^\infty \frac{e^{-\lambda x}(\lambda t)^k}{k!} f_\alpha(t,x)dx \\
&= \frac{(\lambda t^\alpha)^k}{k!} \sum_{j=1}^\infty \frac{(k+j)!}{j!} \frac{(-\lambda t^\alpha)^j}{\Gamma(\alpha(j+k)+1)} = \frac{(\lambda t^\alpha)^k}{k!} E_\alpha^{(k)}(-\lambda t^\alpha) \\
&= (\lambda t^\alpha)^k E_{\alpha, \alpha k + 1}^{k+1}(-\lambda t^\alpha), \quad k = 0, 1, 2..., t \geq 0, \quad 0 < \alpha < 1,
\end{aligned}$$

where f_α is given by (2.3), $E_\alpha(z)$ is the Mittag-Leffler function (2.5), $E_\alpha^{(k)}(z)$ is the $k-$th derivative of $E_\alpha(z)$, and $E^\gamma_{\alpha,\beta}(z)$ is the three-parametric Generalized Mittag-Leffler function defined as follows [17, 40]:

$$E^\gamma_{\alpha,\beta}(z) = \sum_{j=0}^{\infty} \frac{(\gamma)_j z^j}{j!\Gamma(\alpha j+\beta)}, \alpha>0, \beta>0,\ \gamma>0, \quad z\in\mathbb{C}, \tag{3.1}$$

where

$$(\gamma)_j = \begin{cases} 1 & \text{if } j=0; \\ \gamma(\gamma+1)\cdots(\gamma+j-1) & \text{if } j=1,2,\dots \end{cases}$$

is the Pochhammer symbol.

Finally, in [6, 7] it is shown that the marginal distribution of FPP satisfies the following system of fractional differential-difference equations:

$$\mathrm{D}_t^\alpha p_k^{(\alpha)}(t) = -\lambda(p_k^{(\alpha)}(t) - p_{k-1}^{(\alpha)}(t)), \quad k=0,1,2..$$

with initial conditions: $p_0^{(\alpha)}(0)=1, p_k^{(\alpha)}(0)=0, k\geq 1$, and $p_{-1}^{(\alpha)}(t)=0$, where D_t^α is the fractional Caputo-Djrbashian derivative (2.6). See also [13].

Remark. Note that

$$\mathrm{E}N_\alpha(t) = \mathrm{E}\big[\mathrm{E}[N(Y_\alpha(t))|Y_\alpha(t)]\big] = \int_0^\infty [\mathrm{E}N(u)] f_\alpha(t,u)du = \lambda t^\alpha/\Gamma(1+\alpha),$$

where $f_\alpha(t,u)$ is given by (2.3), and [27] showed that

$$\mathrm{Cov}(N_\alpha(t), N_\alpha(s)) = \frac{\lambda(\min(t,s))^\alpha}{\Gamma(1+\alpha)} + \lambda^2 \mathrm{Cov}(Y_\alpha(t), Y_\alpha(s)), \tag{3.2}$$

where $\mathrm{Cov}(Y_\alpha(t), Y_\alpha(s))$ is given in (2.7) while $\mathrm{Cov}(N(t),N(s)) = \lambda\min(t,s)$. In particular,

$$\begin{aligned} \mathrm{Var}N_\alpha(t) &= \lambda^2 t^{2\alpha}\Big[\frac{2}{\Gamma(1+2\alpha)} - \frac{1}{\Gamma^2(1+\alpha)}\Big] + \frac{\lambda t^\alpha}{\Gamma(1+\alpha)} \\ &= \frac{\lambda^2 t^{2\alpha}}{\Gamma^2(1+\alpha)}\Big(\frac{\alpha\Gamma(\alpha)}{\Gamma(2\alpha)} - 1\Big) + \frac{\lambda t^\alpha}{\Gamma(1+\alpha)}, \quad t\geq 0. \end{aligned}$$

The definition of the Hurst index for renewal processes is discussed in [15]. In the same spirit, one can define the analogous of the Hurst index for the FPP as

$$H = \inf\left\{\beta : \limsup_{T\to\infty} \frac{\mathrm{Var}N_\alpha(T)}{T^{2\beta}} < \infty\right\} \in (0,2).$$

To prove the formula (3.2), one can use the conditional covariance formula [42, Exercise 7.20.b]:

$$\mathrm{Cov}(Z_1,Z_2) = \mathrm{E}\big(\mathrm{Cov}(Z_1,Z_2|Y)\big) + \mathrm{Cov}\big(\mathrm{E}(Z_1|Y), \mathrm{E}(Z_2|Y)\big),$$

where Z_1, Z_2 and Y are random variables, and

$$\mathrm{Cov}(Z_1,Z_2|Y) = \mathrm{E}\big((Z_1-\mathrm{E}(Z_1|Y))(Z_2-\mathrm{E}(Z_2|Y))\big).$$

Really, if

$$G_{t,s}(u,v) = \mathrm{P}\{Y_\alpha(t)\leq u, Y_\alpha(s)\leq v\},$$

then $\mathrm{E}(N(Y_\alpha(t))|Y_\alpha(t)) = \mathrm{E}(N(1)) \cdot Y_\alpha(t) = \lambda Y_\alpha(t)$, and

$$\mathrm{Cov}(Y_\alpha(t), Y_\alpha(s)) = \mathrm{Var}\Big(N(1)\int_0^\infty \int_0^\infty \min(u,v) G_{t,s}(du,dv)\Big) + \mathrm{Cov}\big(\lambda Y_\alpha(t), \lambda Y_\alpha(s)\big)$$
$$= \lambda \mathrm{E}(Y_\alpha(\min(t,s))) + \lambda^2 \mathrm{Cov}(Y_\alpha(t), Y_\alpha(s)),$$

since, for example, if $s \le t$, then $v = Y_\alpha(s) \le Y_\alpha(t) = u$, and

$$\int_0^\infty \int_0^\infty v G_{t,s}(du,dv) = \int_0^\infty v \int_0^\infty G_{t,s}(du,dv) = \int_0^\infty v \, d\mathrm{P}\{Y_\alpha(s) \le v\} = \mathrm{E}(Y_\alpha(s)).$$

Remark. For more than one random variable in the condition, the conditional covariance formula becomes more complicated, it can be seen even for the conditional variance formula:

$$\mathrm{Var}(Z) = \mathrm{E}\big(\mathrm{Var}(Z|Y_1,Y_2)\big) + \mathrm{E}\big(\mathrm{Var}[\mathrm{E}(Z|Y_1,Y_2)]|Y_1\big) + \mathrm{Var}\big(\mathrm{E}(Z|Y_1)\big).$$

The corresponding formulas can be found in [10]. That is why for random fields we develop another technique, see Appendix.

3.2. *Watanabe characterization.* Let $(\Omega, \mathcal{F}, \mathrm{P})$ be a complete probability space. Recall that the $\mathcal{F}_t-$adapted, P-integrable stochastic process $M = \{M(t), t \ge 0\}$ is an $\mathcal{F}_t-$martingale (sub-martingale) if $\mathrm{E}(M(t)|\mathcal{F}_s) = (\ge) M(s)$, $0 \le s \le t$, a.s., where $\{\mathcal{F}_t\}$ is a non-decreasing family of sub-sigma fields of $\mathcal{F}$. The following theorem is known as the Watanabe characterization for homogeneous Poisson processes (see, [49] and [12, p. 25]):

Theorem 3.1. *Let $N = \{N(t), t \ge 0\}$ be a $\mathcal{F}_t-$adapted, simple locally finite point process. Then N is a homogeneous Poisson process iff there is a constant $\lambda > 0$, such that the process $M(t) = N(t) - \lambda t$ is an $\mathcal{F}_t-$martingale.*

We extend the well-known Watanabe characterization for FPP.

Theorem 3.2. *Let $X = \{X(t),\, t \ge 0\}$ be a simple locally finite point process. Then X is a FPP iff there exist a constant $\lambda > 0$, and an α-stable subordinator $L_\alpha = \{L_\alpha(t),\, t \ge 0\}$, $0 < \alpha < 1$, such that, denoted by $Y_\alpha(t) = \inf\{s : L_\alpha(s) \ge t\}$ its inverse stable subordinator, the process*

$$M = \{M(t),\, t \ge 0\} = \{X(t) - \lambda Y_\alpha(t),\, t \ge 0\}$$

is a martingale with respect to the induced filtration $\mathcal{F}_t = \sigma(X(s), Y_\alpha(s), s \le t)$.

Proof. If X is a FPP, then $X(t) = N(Y_\alpha(t))$, where Y_α is the inverse of an α-stable subordinator and N is a Poisson process with intensity $\lambda > 0$.

Therefore $N(Y_\alpha(t)) - \lambda Y_\alpha(t)$ is a martingale. Notice that $Y_\alpha(t)$ is continuous increasing and adapted; therefore it is the predictable intensity of the sub-martingale X.

Conversely, it is enough to prove that $X(t) = N(Y_\alpha(t))$, where N is a Poisson process, independent of Y_α.

Consider the inverse of $Y_\alpha(t)$:

$$Z(t) = \inf\{s :\, Y_\alpha(s) \ge t\}.$$

Notice that in general $Z(t)$ does not equal to $L_\alpha(t)$, since for example $Z(t)$ is continuous, but $L_\alpha(t)$ is not.

$\{Z(t),\, t \geq 0\}$ can be seen as a family of stopping times. Then, by the Doob optional sampling theorem [24, Theorem 7.29],

$$M(Z(t)) = X(Z(t)) - \lambda Y_\alpha(Z(t))$$

is still a martingale. Then $X(Z(t)) - \lambda t$ is a martingale. Since $Z(t)$ is increasing, $X(Z(t))$ is a simple point process.

Following the classical Watanabe characterization, $X(Z(t))$ is a classical Poisson process with parameter $\lambda > 0$. Call this process $N(t) = X(Z(t))$. Then $X(t) = N(Y_\alpha(t))$ is a FPP. □

4. Mixed-Fractional Poisson Processes.

4.1. *Definition.* In this section, we consider a more general Mixed-Fractional Poisson process (MFPP)

$$N^{\alpha_1,\alpha_2} = \{N^{\alpha_1,\alpha_2}(t), t \geq 0\} = \{N(Y_{\alpha_1,\alpha_2}(t)), t \geq 0\}, \tag{4.1}$$

where the homogeneous Poisson process N with intensity $\lambda > 0$, and the inverse subordinator Y_{α_1,α_2} given by (2.9) are independent. We will show that N^{α_1,α_2} is the stochastic solution of the system of fractional differential-difference equations: for $k = 0, 1, 2, \ldots$,

$$C_1 \mathrm{D}_t^{\alpha_1} p_k^{(\alpha_1,\alpha_2)}(t) + C_2 \mathrm{D}_t^{\alpha_2} p_k^{(\alpha_1,\alpha_2)}(t) = -\lambda(p_k^{(\alpha_1,\alpha_2)}(t) - p_{k-1}^{(\alpha_1,\alpha_2)}(t)), \tag{4.2}$$

with initial conditions:

$$p_0^{(\alpha_1,\alpha_2)}(0) = 1, p_k^{(\alpha_1,\alpha_2)}(0) = 0, p_{-1}^{(\alpha_1,\alpha_2)}(t) = 0, \quad k \geq 1, \tag{4.3}$$

where D_t^α is the fractional Caputo-Djrbashian derivative (2.6), and for $C_1 \geq 0, C_2 > 0, C_1 + C_2 = 1$, $\alpha_1, \alpha_2 \in (0, 1)$,

$$p_k^{(\alpha_1,\alpha_2)}(t) = \mathrm{P}\{N^{\alpha_1,\alpha_2}(t) = k\}, \qquad k = 0, 1, 2 \ldots$$

4.2. *Distribution Properties.* Using the formulae for Laplace transform of the fractional Caputo-Djrbashian derivative (see, [35, p.39]):

$$\int_0^\infty e^{-st} \mathrm{D}_t^\alpha u(t) dt = s^\alpha \tilde{u}(s) - s^{\alpha-1} u(0), 0 < \alpha < 1,$$

one can obtain from (4.2) with $k = 0$ the following equation

$$C_1 s^{\alpha_1} \tilde{p}_0(s) - C_1 s^{\alpha_1 - 1} + C_2 s^{\alpha 2} \tilde{p}_0(s) - C_2 s^{\alpha_2 - 1} = -\lambda \tilde{p}_0(s), \tilde{p}_0(0) = 1,$$

for the Laplace transform

$$\tilde{p}_0^{(\alpha_1,\alpha_2)}(s) = \tilde{p}_0(s) = \int_0^\infty e^{-st} p_0^{(\alpha_1,\alpha_2)}(t) dt, \quad s \geq 0.$$

Thus

$$\tilde{p}_0(s) = \frac{C_1 s^{\alpha_1 - 1} + C_2 s^{\alpha_2 - 1}}{\lambda + C_1 s^{\alpha_1} + C_2 s^{\alpha_2}}, \quad s \geq 0,$$

and using the formula for an inverse Laplace transform (see, [17]), for $\Re\alpha > 0, \Re\beta > 0, \Re s > 0, \Re(\alpha - \rho) > 0, \Re(\alpha - \beta) > 0$, and $|as^\beta/(s^\alpha + b)| < 1$:

$$\mathcal{L}^{-1}\Big(\frac{s^{\rho-1}}{s^\alpha + as^\beta + b}; t\Big) = t^{\alpha-\rho} \sum_{r=0}^{\infty} (-a)^r t^{(\alpha-\beta)r} E^{r+1}_{\alpha,\alpha+(\alpha-\beta)r-\rho+1}(-bt^\alpha), \tag{4.4}$$

one can find an exact form of the $p_0^{(\alpha_1,\alpha_2)}(t)$ in terms of generalized Mittag-Leffler functions (3.1):

$$p_0^{(\alpha_1,\alpha_2)}(t) = \sum_{r=0}^{\infty}\left(-\frac{C_1}{C_2}t^{\alpha_2-\alpha_1}\right)^r E^{r+1}_{\alpha_2,(\alpha_2-\alpha_1)r+1}\left(-\frac{\lambda}{C_2}t^{\alpha_2}\right) \tag{4.5}$$
$$-\sum_{r=0}^{\infty}\left(-\frac{C_1}{C_2}t^{\alpha_2-\alpha_1}\right)^{r+1} E^{r+1}_{\alpha_2,(\alpha_2-\alpha_1)(r+1)+1}\left(-\frac{\lambda}{C_2}t^{\alpha_2}\right).$$

For $k \geq 1$,we obtain from (4.2):

$$\tilde{p}_k(s)(\lambda + C_1 s^{\alpha_1} + C_2 s^{\alpha_2}) = \lambda \tilde{p}_{k-1}(s),$$

where

$$\tilde{p}_k^{(\alpha_1,\alpha_2)}(s) = \tilde{p}_k(s) = \int_0^{\infty} e^{-st} p_k^{(\alpha_1,\alpha_2)}(t)dt, \quad s \geq 0.$$

Thus from (4.2) we obtain the following expression for the Laplace transform of $p_k^{(\alpha_1,\alpha_2)}(t)$, $k \geq 0$:

$$\tilde{p}_k(s) = \left(\frac{\lambda}{\lambda + C_1 s^{\alpha_1} + C_2 s^{\alpha_2}}\right)\tilde{p}_{k-1}(s) = \left(\frac{\lambda}{\lambda + C_1 s^{\alpha_1} + C_2 s^{\alpha_2}}\right)^k \tilde{p}_0(s) \tag{4.6}$$
$$= \frac{\lambda^k (C_1 s^{\alpha_1 - 1} + C_2 s^{\alpha_2 - 1})}{(\lambda + C_1 s^{\alpha_1} + C_2 s^{\alpha_2})^{k+1}} = \frac{\lambda^k (C_1 s^{\alpha_1} + C_2 s^{\alpha_2})}{s(\lambda + C_1 s^{\alpha_1} + C_2 s^{\alpha_2})^{k+1}}, \ k = 0, 1, 2...$$

On the other hand, one can compute the Laplace transform from the stochastic representation (4.1). If

$$p_k^{(\alpha_1,\alpha_2)}(t) = \mathrm{P}\{N(Y_{\alpha_1,\alpha_2}(t) = k\} = \int_0^{\infty} \frac{e^{-\lambda x}}{k!}(\lambda x)^k f_{\alpha_1,\alpha_2}(t,x)dx, \tag{4.7}$$

where $f_{\alpha_1,\alpha_2}(t,x)$ is given by (2.13), then using (2.11),(2.12) we have for $k \geq 0, s > 0$

$$\tilde{p}_k(s) = \int_0^{\infty} e^{-st} p_k^{(\alpha_1,\alpha_2)}(t)dt = \int_0^{\infty} \frac{e^{-\lambda x}}{k!}(\lambda x)^k [\int_0^{\infty} e^{-st} f_{\alpha_1,\alpha_2}(t,x)dt]dx$$
$$= \frac{\lambda^k}{k!}\frac{\phi(s)}{s}\int_0^{\infty} e^{-\lambda x} x^k e^{-x\phi(s)}dx$$

Note that

$$\frac{\partial^k}{\partial \lambda^k}\int_0^{\infty} e^{-\lambda x} e^{-x\phi(s)}dx = (-1)^k \int_0^{\infty} e^{-\lambda x} x^k e^{-x\phi(s)}dx$$
$$= \frac{\partial^k}{\partial \lambda^k}\frac{1}{\lambda + \phi(s)} = (-1)^k \frac{k!}{(\lambda + \phi(s))^{k+1}};$$

thus

$$\tilde{p}_k(s) = \lambda^k \frac{\phi(s)}{s(\lambda + \phi(s))^{k+1}} = \frac{\lambda^k (C_1 s^{\alpha_1} + C_2 s^{\alpha_2})}{s(\lambda + C_1 s^{\alpha_1} + C_2 s^{\alpha_2})^{k+1}},$$

the same expression as (4.6). We can formulate the result in the following form:

THEOREM 4.1. *The MFPP N^{α_1,α_2} defined in (4.1) is the stochastic solution of the system of fractional differential-difference equations (4.2) with initial conditions (4.3).*

Note that in [5] one can find some other stochastic representations of the MFPP (4.1). Also, some analytical expression for $p_0^{(\alpha_1,\alpha_2)}(t)$ is given by (4.5), while the analytical expression for $p_k^{(\alpha_1,\alpha_2)}(t)$,for $k \geq 1$, are given by (4.7).

Moreover, $p_k^{(\alpha_1,\alpha_2)}(t)$,for $k \geq 1$, can be obtained by the following recurrent relation:

$$p_k^{(\alpha_1,\alpha_2)}(t) = \int_0^t p_{k-1}^{(\alpha_1,\alpha_2)}(t-z)g(z)dz,$$

where

$$\tilde{g}(s) = \int_0^\infty e^{-st}g(z)dz = \frac{\lambda}{\lambda + C_1 s^{\alpha_1} + C_2 s^{\alpha_2}},$$

and from (4.4):

$$g(z) = \frac{\lambda}{C_2} z^{\alpha_2 - 1} \sum_{r=0}^{\infty} \Big(-\frac{C_1}{C_2} t^{\alpha_2-\alpha_1} \Big)^r E^{r+1}_{\alpha_2,\alpha_2+(\alpha_2-\alpha_1)r}\Big(-\frac{\lambda}{C_2} t^{\alpha_2} \Big).$$

4.3. *Dependence.* From [27, Theorem 2.1] and (2.10), we have the following expressions for moments in form of the function

$$\begin{aligned}
U(t) &= \frac{1}{C_2} t^{\alpha_2} E_{\alpha_2-\alpha_1,\alpha_2+1}(-C_1 t^{\alpha_2-\alpha_1}/C_2),\\
\mathrm{E}N^{\alpha_1,\alpha_2}(t) &= \lambda U(t),\\
\mathrm{Var}N^{\alpha_1,\alpha_2}(t) &= \lambda^2 \frac{1}{C_2^2} t^{2\alpha_2} [2E_{\alpha_2-\alpha_1,\alpha_1+\alpha_2+1}(-C_1 t^{\alpha_2-\alpha_1}/C_2)\\
&\quad - (E_{\alpha_2-\alpha_1,\alpha_2+1}(-C_1 t^{\alpha_2-\alpha_1}/C_2))^2]\\
&\quad + \lambda \frac{1}{C_2} t^{\alpha_2} E_{\alpha_2-\alpha_1,\alpha_2+1}(-C_1 t^{\alpha_2-\alpha_1}/C_2),\\
\mathrm{Cov}(N^{\alpha_1,\alpha_2}(t), N^{\alpha_1,\alpha_2}(s)) &= \lambda U(\min(t,s)) + \lambda^2 \Big\{ \int_0^{\min(t,s)} \Big(U(t_1-\tau)\\
&\quad + U(t_2-\tau)\Big) dU(\tau) - U(t_1)U(t_2) \Big\}.
\end{aligned}$$

We extend the Watanabe characterization for MFPP. Let $\Lambda(t) : \mathbb{R}_+ \to \mathbb{R}_+$ be a non-negative right-continuous non-decreasing deterministic function such that $\Lambda(0) = 0$, $\Lambda(\infty) = \infty$, and $\Lambda(t) - \Lambda(t-) \leq 1$ for any t. Such a function will be called *consistent*. The Mixed-Fractional Non-homogeneous Poisson process (MFNPP) is defined as

$$N_\Lambda^{\alpha_1,\alpha_2} = \{N_\Lambda^{\alpha_1,\alpha_2}(t), t \geq 0\} = \{N(\Lambda(Y_{\alpha_1,\alpha_2}(t))), t \geq 0\},$$

where the homogeneous Poisson process N with intensity $\lambda = 1$, and the inverse subordinator Y_{α_1,α_2} given by (2.9) are independent.

THEOREM 4.2. *Let $X = \{X(t), t \geq 0\}$ be a simple locally finite point process. X is a MFNPP iff there exist a consistent function $\Lambda(t)$, and a mixed stable subordinator $\{L_{\alpha_1,\alpha_2}(t), t \geq 0\}$, $0 < \alpha_1 < 1$, $0 < \alpha_2 < 1$, defined in (2.8), such that*

$$M = \{M(t), t \geq 0\} = \{X(t) - \Lambda(Y_{\alpha_1,\alpha_2}(t)), t \geq 0\}$$

is a martingale with respect to the induced filtration $\mathcal{F}_t = \sigma(X(s), Y_{\alpha_1,\alpha_2}(s), s \leq t)$, where $Y_{\alpha_1,\alpha_2}(t) = \inf\{s : L_{\alpha_1,\alpha_2}(t) \geq t\}$ is the inverse mixed stable subordinator.

PROOF. If X is a MFNPP, then $X(t) = N(\Lambda(Y_{\alpha_1,\alpha_2}(t)))$, where the homogeneous Poisson process N and the inverse subordinator Y_{α_1,α_2} given by (2.9) are independent. Therefore $X(t)-\Lambda(Y_{\alpha_1,\alpha_2}(t))$ is a martingale.

Conversely, it is enough to prove that $X(t) = N(\Lambda(Y_{\alpha_1,\alpha_2}(t)))$, where N is a Poisson process, independent of Y_α.

Consider the pseudo inverse of $Y_{\alpha_1,\alpha_2}(t)$:

$$Z(t) = \inf\{s : Y_{\alpha_1,\alpha_2}(s) \geq t\}.$$

$\{Z(t), t \geq 0\}$ can be seen as a family of stopping times. Then, by the Doob optional stopping theorem,

$$M(Z(t)) = X(Z(t)) - \Lambda(Y_{\alpha_1,\alpha_2}(Z(t)))$$

is still a martingale. Then $X(Z(t)) - \Lambda(t)$ is a martingale. Since $Z(t)$ is increasing, $X(Z(t))$ is a simple point process.

Following the extension of Watanabe characterization given in [11], $X(Z(t))$ is a non-homogeneous Poisson process with deterministic intensity $\Lambda(t)$. Call this process $N(t) = X(Z(t))$. Then $X(t) = \Lambda(Y_{\alpha_1,\alpha_2}(Z(t)))$ is a MFNPP. □

5. Two-Parameter Fractional Poisson Processes and Martingales.

5.1. *Homogeneous Poisson random fields.* This section collects some known results from the theory of two-parameter Poisson processes and homogeneous Poisson random fields (PRF) (see, e.g., [43, 36], among the others).

Let $(\Omega, \mathcal{F}, P)$ be a complete probability space and let $\left\{\mathcal{F}_{t_1,t_2}; (t_1,t_2) \in \mathbb{R}^2_+\right\}$ be a family of sub-σ-fields of $\mathcal{F}$ such that

(i) $\mathcal{F}_{s_1,s_2} \subseteq \mathcal{F}_{t_1,t_2}$ for any $\quad s_1 \leq t_1, \quad s_2 \leq t_2$;

(ii) $\mathcal{F}_{0,0}$ contains all null sets of $\mathcal{F}$;

(iii) for each $z \in \mathbb{R}^2_+$, $\mathcal{F}_z = \bigcap_{z \prec z'} \mathcal{F}_{z'}$ where $z = (s_1,s_2) \prec z' = (t_1,t_2)$ denotes the partial order on $\mathbb{R}^2_+$, which means that $s_1 \leq t_1, \quad s_2 \leq t_2$.

Given $(s_1,s_2) \prec (t_1,t_2)$ we denote by

$$\Delta_{s_1,s_2}X(t_1,t_2) = X(t_1,t_2) - X(t_1,s_2) - X(s_1,t_2) + X(s_1,s_2)$$

the increments of the random field $X(t_1,t_2), (t_1,t_2) \in \mathbb{R}^2_+$ over the rectangle $((s_1,s_2),(t_1,t_2)]$. In addition, we denote

$$\mathcal{F}_{\infty,t_2} = \sigma(\mathcal{F}_{t_1,t_2}, t_1 > 0), \mathcal{F}_{t_1,\infty} = \sigma(\mathcal{F}_{t_1,t_2}, t_2 > 0), \text{ and } \mathcal{F}_{\infty,s_2} \vee \mathcal{F}_{s_1,\infty} = \sigma(\mathcal{F}_{t_1,\infty}, \mathcal{F}_{\infty,t_2}).$$

A strong martingale is an integrable two-parameter process X such that

$$E(\Delta_{s_1,s_2}X(t_1,t_2)|\mathcal{F}_{\infty,s_2} \vee \mathcal{F}_{s_1,\infty}) = 0,$$

for any $z = (s_1,s_2) \prec z' = (t_1,t_2) \in \mathbb{R}^2_+$.

Let $\{\mathcal{F}_{t_1,t_2}\}$ be a family of sub-σ-fields of $\mathcal{F}$ satisfying the previous conditions (i), (ii), (iii) for all $(t_1,t_2) \in \mathbb{R}^2_+$. An $\mathcal{F}_{t_1,t_2}-$ PRF is adapted, a cadlag field $N = \left\{N(t_1,t_2), (t_1,t_2) \in \mathbb{R}^2_+\right\}$, such that,

(1) $N(t_1,0) = N(0,t_2) = 0$ a.s.

(2) for all $(s_1,s_2) \prec (t_1,t_2)$ the increments $\Delta_{s_1,s_2}N(t_1,t_2)$ are independent of $\mathcal{F}_{\infty,s_2} \vee \mathcal{F}_{s_1,\infty}$, and has a Poisson law with parameter $\lambda\,(t_1-s_1)\,(t_2-s_2)$, that is,

$$\mathrm{P}\left\{\Delta_{s_1,s_2}N(t_1,t_2) = k\right\} = \frac{e^{-\lambda|S|}\left(\lambda\,|S|\right)^k}{k!}, \quad \lambda > 0, \quad k = 0,1,\ldots,$$

where $S = ((s_1, s_2), (t_1, t_2)]$, $\lambda > 0$, and $|S|$ is the Lebesgue measure of S.

If we do not specify the filtration, $\{\mathcal{F}_{t_1,t_2}\}$ will be the filtration generated by the field itself, completed with the nulls sets of $\mathcal{F}^N = \sigma\left\{N(t_1,t_2), (t_1,t_2) \in \mathbb{R}^2_+\right\}$.

It is known that then there is a simple locally finite point random measure $N(\cdot)$, such that for any finite $n = 1, 2, \ldots$, and for any disjoint bounded Borel sets $A_1, ..., A_n$

$$\begin{aligned}&\mathrm{P}\left(N(A_1) = k_1, ..., N(A_n) = k_n\right)\\ &\quad = \frac{\lambda^{k_1+...+k_n}}{k_1! \cdot .. \cdot k_n!}(|A_1|)^{k_1} \cdots (|A_n|)^{k_n} \exp\left\{-\sum_{j=1}^{n} \lambda |A_j|\right\}, \quad k_j = 0, 1, 2, ...,\end{aligned}$$

while

$$\mathrm{E}N(A) = \lambda |A|, \ \mathrm{Cov}(N(A_1), N(A_2)) = \lambda |A_1 \cap A_2|.$$

THEOREM 5.1 (Two Parameter Watanabe Theorem [20]). *A random simple locally finite counting measure N is a two-parameter PRF iff $N(t_1, t_2) - \lambda t_1 t_2$ is a strong martingale.*

5.2. *Fractional Poisson random fields.* Let $Y^{(1)}_{\alpha_1}(t), t \geq 0$ and $Y^{(2)}_{\alpha_2}(t), t \geq 0$ be two independent inverse stable subordinators with indices $\alpha_1 \in (0, 1)$ and $\alpha_2 \in (0, 1)$, which are independent of the Poisson field $N(t_1, t_2), (t_1, t_2) \in \mathbb{R}^2_+$. In [26] it is defined the Fractional Poisson field (FPRF) as follows:

$$N_{\alpha_1,\alpha_2}(t_1, t_2) = N(Y^{(1)}_{\alpha_1}(t_1), Y^{(2)}_{\alpha_2}(t_2)), \quad (t_1, t_2) \in \mathbb{R}^2_+. \tag{5.1}$$

We obtain the marginal distribution of FPRF: for $k = 0, 1, 2, \ldots$,

$$\begin{aligned}p^{\alpha_1,\alpha_2}_k(t_1, t_2) &= \mathrm{P}\left(N_{\alpha_1,\alpha_2}(t_1, t_2) = k\right)\\ &= \int_0^\infty \int_0^\infty \frac{e^{-\lambda x_1 x_2}(\lambda x_1 x_2)^k}{k!} f_{\alpha_1}(t_1, x_1) f_{\alpha_2}(t_2, x_2) dx_1 dx_2,\end{aligned} \tag{5.2}$$

where $f_a(t, x)$ is given by (2.3). In other words, for $(t_1, t_2) \in \mathbb{R}^2_+$, $k = 0, 1, \ldots$

$$\begin{aligned}\mathrm{P}\left(N_{\alpha_1,\alpha_2}(t_1, t_2) = k\right) &= \frac{t_1 t_2 \lambda^k}{\alpha_1 \alpha_2 k!} \int_0^\infty \int_0^\infty e^{-\lambda x_1 x_2} x_1^{k-1-\frac{1}{\alpha_1}} x_2^{k-1-\frac{1}{\alpha_2}} g_{\alpha_1}(t_1 x_1^{-\frac{1}{\alpha_1}}) g_{\alpha_2}(t_2 x_2^{-\frac{1}{\alpha_2}}) dx_1 dx_2,\\ &= \frac{\lambda^k}{k! t_1 t_2} \int_0^\infty \int_0^\infty e^{-\lambda x_1 x_2} x_1^{k+\frac{1}{\alpha_1}} x_2^{k+\frac{1}{\alpha_2}} W_{-\alpha_1,0}(-\frac{x_1}{t_1^{\alpha_1}}) W_{-\alpha_2,0}(-\frac{x_2}{t_2^{\alpha_2}}) dx_1 dx_2,\end{aligned} \tag{5.3}$$

where the Wright generalized Bessel function is defined by (2.2), and $g_\alpha(x)$ is defined by (2.1).

Using the Laplace transform given by (2.4) one can obtain an exact expression for the double Laplace transform of (5.2): for $k = 0, 1, 2, \ldots$,

$$\begin{aligned}\mathcal{L}\{p_k(t_1, t_2); s_1, s_2\} &= \int_0^\infty \int_0^\infty e^{-s_t t_1 - s_2 t_2} p_k(t_1, t_2) dt_1 dt_2\\ &= \int_0^\infty \int_0^\infty \frac{e^{-\lambda x_1 x_2}(\lambda x_1 x_2)^k}{k!} s_1^{\alpha_1 - 1} s_2^{\alpha_2 - 1} \exp\{-x_1 s_1^{\alpha_1} - x_2 s_2^{\alpha_2}\} dx_1 dx_2.\end{aligned} \tag{5.4}$$

It is known (see, e.g., [4, Theorems 1.1. and 1.2]) that a single point random field M can be regarded as a random set of points and could be characterized by its system of finite-dimensional distributions as

$$\mathrm{P}\left(M(B_1) = k_1, ..., M(B_m) = k_m\right)$$

for all integers $m > 0$ and all compacts $B_1, .., B_m$.

Note that

$$\begin{aligned} \mathrm{E}N_{\alpha_1,\alpha_2}(t_1,t_2) &= \mathrm{E}\big[\mathrm{E}[N(Y_{\alpha_1}(t_1), Y_{\alpha_2}(t_2))|Y_{\alpha_1}(t_1), Y_{\alpha_2}(t_2)]\big] \\ &= \int_0^\infty \int_0^\infty \mathrm{E}N(u_1,u_2) f_{\alpha_1}(t_1,u_1) f_{\alpha_2}(t_2,u_2) du_1 du_2 \\ &= \lambda t_1^{\alpha_1} t_2^{\alpha_2} / [\Gamma(1+\alpha_1)\Gamma(1+\alpha_2)] \end{aligned} \tag{5.5}$$

and, for $(t_1,t_2),(s_1,s_2) \in \mathbb{R}^2_+$,

$$\begin{aligned} &\mathrm{Cov}(N_{\alpha_1,\alpha_2}(t_1,t_2), N_{\alpha_1,\alpha_2}(s_1,s_2)) \\ &= \lambda^2 \Big\{ \Big[\frac{1}{\Gamma(1+\alpha_1)\Gamma(\alpha_1)} \int_0^{\min(t_1,s_1)} (t_1-\tau_1)^{\alpha_1} + (s_1-\tau_1)^{\alpha_1}) \tau_1^{\alpha_1-1} d\tau_1 - \frac{(s_1 t_1)^{\alpha_1}}{\Gamma^2(1+\alpha_1)} \Big] \\ &\quad \times \Big[\frac{1}{\Gamma(1+\alpha_2)\Gamma(\alpha_2)} \int_0^{\min(t_2,s_2)} (t_2-\tau_2)^{\alpha_2} + (s_2-\tau_2)^{\alpha_2}) \tau_2^{\alpha_2-1} d\tau_2 - \frac{(s_2 t_2)^{\alpha_2}}{\Gamma^2(1+\alpha_2)} \Big] \\ &\quad + \Big[\frac{(s_1 t_1)^{\alpha_1}}{\Gamma^2(1+\alpha_1)} - \frac{2}{\Gamma(2+\alpha_1)} (\min(t_1,s_1))^{1+\alpha_1} \Big] \\ &\quad \times \Big[\frac{1}{\Gamma(1+\alpha_2)\Gamma(\alpha_2)} \int_0^{\min(t_2,s_2)} (t_2-\tau_2)^{\alpha_2} + (s_2-\tau_2)^{\alpha_2}) \tau_2^{\alpha_2-1} d\tau_2 - \\ &\qquad \frac{2}{\Gamma(2+\alpha_2)} (\min(t_2,s_2))^{1+\alpha_2} \Big] + \Big[\frac{(s_2 t_2)^{\alpha_2}}{\Gamma^2(1+\alpha_2)} - \frac{2}{\Gamma(2+\alpha_2)} (\min(t_2,s_2))^{1+\alpha_2} \Big] \\ &\quad \times \Big[\frac{1}{\Gamma(1+\alpha_1)\Gamma(\alpha_1)} \int_0^{\min(t_1,s_1)} (t_1-\tau_1)^{\alpha_1} + (s_1-\tau_1)^{\alpha_1}) \tau_1^{\alpha_1-1} d\tau_1 - \\ &\qquad \frac{2}{\Gamma(2+\alpha_1)} (\min(t_1,s_1))^{1+\alpha_1} \Big] \Big\} + \lambda \frac{(\min(t_1,s_1))^{\alpha_1} (\min(t_2,s_2))^{\alpha_2}}{\Gamma(1+\alpha_1)\Gamma(1+\alpha_2)}; \end{aligned} \tag{5.6}$$

in particular, for $(t_1,t_2),(s_1,s_2) \in \mathbb{R}^2_+$,

$$\mathrm{Var}N_{\alpha_1,\alpha_2}(t_1,t_2) = \lambda^2 \{ {t_1}^{\alpha_1+1} {t_2}^{\alpha_2+1} C_1(\alpha_1,\alpha_2) - \lambda^2 {t_1}^{2\alpha_1} {t_2}^{2\alpha_2} C_2(\alpha_1,\alpha_2) + \lambda {t_1}^{\alpha_1} {t_2}^{\alpha_2} C_3(\alpha_1,\alpha_2) \}, \tag{5.7}$$

where

$$\begin{aligned} C_1(\alpha_1,\alpha_2) &= \frac{4}{\Gamma(2+\alpha_1)\Gamma(2+\alpha_2)}; \quad C_2(\alpha_1,\alpha_2) = \frac{1}{\Gamma^2(1+\alpha_1)\Gamma^2(1+\alpha_2)}; \\ C_3(\alpha_1,\alpha_2) &= \frac{1}{\Gamma(1+\alpha_1)\Gamma(1+\alpha_2)}. \end{aligned}$$

We can summarize our results in the following

PROPOSITION 5.1. *Let $N_{\alpha_1,\alpha_2}(t_1,t_2),(t_1,t_2) \in \mathbb{R}^2_+$, be a FPRF defined by (5.1). Then*

i) $\mathrm{P}\left(N_{\alpha_1,\alpha_2}(t_1,t_2) = k\right)$, $k = 0,1,2...$ *is given by (5.3);*

ii) $\mathrm{E}N_{\alpha_1,\alpha_2}(t_1,t_2), \mathrm{Var}N_{\alpha_1,\alpha_2}(t_1,t_2)$ *and* $\mathrm{Cov}(N_{\alpha_1,\alpha_2}(t_1,t_2), N_{\alpha_1,\alpha_2}(s_1,s_2))$ *are given by (5.5), (5.7), (5.6), respectively.*

The proof of the theorem is given in the Appendix.

REMARK. Note that formulae (5.5),(5.7) and (5.6) are corrected versions of the formulae (26) and (27) of the paper [26].

REMARK. Following the ideas of this paper, the Hurst index of the Fractional Poisson random field in $d = 2$ can be defined as follows:

$$H = \inf\left\{\beta : \limsup_{T\to\infty} \frac{\mathrm{Var}N_{\alpha_1,\alpha_2}(T,T)}{T^{2d\beta}} < \infty\right\} \in (0,4).$$

REMARK. Any random field

$$Z(t_1,t_2) = N(Y_1(t_1), Y_2(t_2)), \quad (t_1,t_2) \in \mathbb{R}^2_+$$

defined on the positive quadrant $\mathbb{R}^2_+$ can be extended in the whole space $\mathbb{R}^2$ in the following way: let $Z_j(t_1,t_2), (t_1,t_2) \in \mathbb{R}^2_+,\ j = 1,2,3,4$ be independent copies of the random field $Z(t_1,t_2), (t_1,t_2) \in \mathbb{R}^2_+$.

Then one can define

$$\bar{Z}(t_1,t_2) = \begin{cases} Z_1(t_1,t_2), & t_1 \geq 0, t_2 \geq 0 \\ -Z_2(-t_1^-,-t_2), & t_1 < 0, t_2 \geq 0 \\ -Z_3(t_1,-t_2^-), & t_1 \geq 0, t_2 < 0 \\ Z_4(-t_1^-,-t_2^-), & t_1 < 0, t_2 < 0 \end{cases}$$

Therefore, modifying the cadlag property we obtain a Poisson like random field $\bar{Z}(t_1,t_2), (t_1,t_2) \in \mathbb{R}^2$ which has a similar covariance structure (replacing t_1, t_2, s_1, s_2 by $|t_1|, |t_2|, |s_1|, |s_2|$).

5.3. *Characterization on increasing paths.* Let $L_\alpha = \{L_\alpha(t), t \geq 0\}$, be an α-stable subordinator, and $Y_\alpha = \{Y_\alpha(t), t \geq 0\}$ be its inverse ($\alpha \in (0,1)$). Recall that $L_\alpha(t)$ is a cadlag strictly increasing process t, while $Y_\alpha\ (t)$ is nondecreasing and continuous. As a consequence, the latter defines a random nonnegative measure μ_α on $(\mathbb{R}_+, \mathcal{B}_{\mathbb{R}_+})$ such that $\mu_\alpha([0,t]) = Y_\alpha(t)$. The σ-algebra $\mathcal{G}$ contains all the information given by μ_α:

$$\mathcal{G} := \sigma(L_\alpha(t), t \geq 0) = \sigma(Y_\alpha(t), t \geq 0) = \sigma(\mu_\alpha(B), B \in \mathcal{B}_{\mathbb{R}_+}).$$

Now, let $X(t) = N(Y_\alpha(t))$ be a FPP, where N has intensity λ. We denote by $\{\mathcal{F}_t^X, t \in \mathbb{R}_+\}$ its natural filtration. We note that each $\mu_\alpha([0,t])$ is $\mathcal{G}$-measurable, while $N(w) - N(\mu_\alpha([0,s]))$ is independent of $\sigma(\mathcal{F}_s^X, \mathcal{G})$ for any $w \geq \mu_\alpha([0,s])$. Hence, for any bounded $\mathcal{F}_s^X$-measurable random variable $Y(s)$, we have

$$\begin{aligned} \mathrm{E}\Big(\int_0^\infty Y(s)\mathbb{1}_{(s,t]}dX_v\Big) &= \mathrm{E}\Big(Y(s)\mathrm{E}\Big(\int_0^\infty \mathbb{1}_{(s,t]}N(\mu_\alpha(dv))\Big|\sigma(\mathcal{F}_s^X,\mathcal{G})\Big)\Big) \\ &= \mathrm{E}\Big(Y(s)\int_0^\infty \mathbb{1}_{(\mu_\alpha([0,s]),\mu_\alpha([0,t])]}\mathrm{E}(dN_w|\sigma(\mathcal{F}_s^X,\mathcal{G}))\Big) \\ &= \mathrm{E}\Big(Y(s)\lambda\mu_\alpha([s,t])\Big) \\ &= \mathrm{E}\Big(\int_0^\infty Y(s)\mathbb{1}_{(s,t]}\lambda\mu_\alpha(dv)\Big). \end{aligned}$$

In other words, by [12, Theorem T4], the FPP X is a doubly stochastic Poisson process with respect to the filtration $\{\sigma(\mathcal{F}_t^X, \mathcal{G}), t \in \mathbb{R}_+\}$. Therefore a first characterization of a FPP may be written in the following way.

COROLLARY 5.1. *A process N_α is a FPP iff it is a doubly stochastic Poisson process with intensity λY_α, with respect to the filtration $\{\sigma(\mathcal{F}_t^X, \mathcal{G}), t \in \mathbb{R}_+\}$. In other words, whenever $B_1, \ldots, B_n$ are disjoint bounded Borel sets and $x_1, \ldots, x_n$ are non-negative integers, then*

$$\mathrm{P}\Big(\bigcap_{i=1}^n \{N_\alpha(B_i) = x_i\} \Big| \mathcal{G}\Big) = \prod_{i=1}^n \frac{\exp(-\lambda\mu_\alpha(B_i))(\lambda\mu_\alpha(B_i))^{x_i}}{x_i!}.$$

An analogous result may be found for FPRF. In fact, let $Y_{\alpha_1}^{(1)}(t), t \geq 0$ and $Y_{\alpha_2}^{(2)}(t), t \geq 0$ be two independent inverse stable subordinators with indices $\alpha_1 \in (0,1)$ and $\alpha_2 \in (0,1)$. Let μ_{α_1} and μ_{α_2}, $\mathcal{G}_1$ and $\mathcal{G}_2$ as above.

If $\mu_{\boldsymbol{\alpha}} = \mu_{\alpha_1} \otimes \mu_{\alpha_2}$ is the product measure and $\mathcal{G} = \sigma(\mathcal{G}_1, \mathcal{G}_2)$, we can mimic the same reasoning as above once we have noted that $\Delta_{\mu_{\alpha_1}([0,s_1]),\mu_{\alpha_2}([0,s_2])} N(w_1, w_2)$ and $\sigma(\mathcal{F}^X_{\infty,s_2} \vee \mathcal{F}^X_{s_1,\infty}))$ are conditionally independent, given , $\mathcal{G}$. In fact

$$\begin{aligned}
&\mathrm{E}\Big(\Delta_{s_1,s_2} X(t_1,t_2) \Big| \sigma(\mathcal{F}^X_{\infty,s_2} \vee \mathcal{F}^X_{s_1,\infty}, \mathcal{G}))\Big) \\
&\qquad = \mathrm{E}\Big(\Delta_{\mu_{\alpha_1}([0,s_1]),\mu_{\alpha_2}([0,s_2])} N(\mu_{\alpha_1}([0,t_1]), \mu_{\alpha_2}([0,t_2])) \Big| \sigma(\mathcal{F}^X_{\infty,s_2} \vee \mathcal{F}^X_{s_1,\infty}, \mathcal{G}))\Big) \\
&\qquad = \mathrm{E}\Big(\Delta_{\mu_{\alpha_1}([0,s_1]),\mu_{\alpha_2}([0,s_2])} N(\mu_{\alpha_1}([0,t_1]), \mu_{\alpha_2}([0,t_2])) | \mathcal{G}) \\
&\qquad = \lambda\mu_{\boldsymbol{\alpha}}(((s_1,s_2),(t_1,t_2)]).
\end{aligned} \tag{5.8}$$

In other words, with the notation given in [36], the FPRF X is a $\mathcal{F}^*$-doubly stochastic Poisson process with respect to the filtration $\{\sigma(\mathcal{F}^X_{t_1,t_2}, \mathcal{G})), t_1, t_2 \in \mathbb{R}_+\}$ by [36, Theorem 1]. Again, we may summarize this result in the following statement.

PROPOSITION 5.2. *A process N_{α_1,α_2} is a FPRF iff it is a $\mathcal{F}^*$-doubly stochastic Poisson process with intensity $\lambda Y_{\alpha_1} \cdot Y_{\alpha_2}$, with respect to the filtration $\{\sigma(\mathcal{F}^X_{t_1,t_2}, \mathcal{G})), t_1, t_2 \in \mathbb{R}_+\}$. In other words, whenever $B_1, \ldots, B_n$ are disjoint bounded Borel sets in $\mathbb{R}_+ \times \mathbb{R}_+$ and $x_1, \ldots, x_n$ are non-negative integers, then*

$$\mathrm{P}\Big(\bigcap_{i=1}^n \{N_{\alpha_1,\alpha_2}(B_i) = x_i\} \Big| \mathcal{G}\Big) = \prod_{i=1}^n \frac{\exp(-\lambda\mu_{\boldsymbol{\alpha}}(B_i))(\lambda\mu_{\boldsymbol{\alpha}}(B_i))^{x_i}}{x_i!}. \tag{5.9}$$

Now, let $t_1 > 0$ be fixed. The process $t \mapsto N_{\alpha_1,\alpha_2}(t_1, t)$ is the trace of the FPRF along the increasing t-indexed family of sets $t \mapsto [(0,0),(t_1,t)]$. As a consequence of the previous results, we obtain:

THEOREM 5.2. *A random simple locally finite counting measure N_{α_1,α_2} is a FPRF iff $\mathcal{G}_1, \mathcal{G}_2$ are independent, and fixed $t_1, t_2 \geq 0$, the process $N_{\alpha_1,\alpha_2}(t_1, t)$, conditioned on $\mathcal{G}_1$, is a FPP $N_{\alpha_2}(t)$, the process $N_{\alpha_1,\alpha_2}(t, t_2)$, conditioned on $\mathcal{G}_2$, is a FPP $N_{\alpha_1}(t)$, and the two processes $N_{\alpha_1}(t_1 + t) - N_{\alpha_1}(t_1), N_{\alpha_2}(t_2 + t) - N_{\alpha_1}(t_2)$ are conditionally independent given $\sigma(\mathcal{G}, \sigma(N_{\alpha_1,\alpha_2}(s_1, s_2), (s_1, 2_2) \prec (t_1, t_2)))$.*

PROOF. Assume that N_{α_1,α_2} is a FPRF and $t_1 > 0$ fixed. Denote by $X_t = N_{\alpha_1,\alpha_2}(t_1, t)$ and note that $\sigma(\{Y_{\alpha_2}(t), t \geq 0\}) = \mathcal{G}_2$. Let $B_1, \ldots, B_n$ be disjoint bounded Borel sets and $x_1, \ldots, x_n$

non-negative integers. We have

$$\begin{aligned}\mathrm{P}\Big(\bigcap_{i=1}^{n}\{N_{\alpha_1,\alpha_2}([0,t_1]\times B_i)=x_i\}\Big|\sigma(\mathcal{G}_1,\sigma(\{Y_{\alpha_2}(t),t\geq 0\}))\Big)&=\mathrm{P}\Big(\bigcap_{i=1}^{n}\{N_{\alpha_1,\alpha_2}([0,t_1]\times B_i)=x_i\}\Big|\mathcal{G}\Big)\\&=\prod_{i=1}^{n}\frac{\exp(-\lambda\mu_{\boldsymbol{\alpha}}([0,t_1]\times B_i))(\lambda\mu_{\boldsymbol{\alpha}}([0,t_1]\times B_i))^{x_i}}{x_i!}\\&=\prod_{i=1}^{n}\frac{\exp(-\lambda Y_{\alpha_1}(t_1)\cdot\mu_{\alpha_2}(B_i))(\lambda Y_{\alpha_1}(t_1)\cdot\mu_{\alpha_2}(B_i))^{x_i}}{x_i!},\end{aligned}$$

and hence $X_t = M(Y_{\alpha_2}(t))$, where, conditioned on $\mathcal{G}_1$, M is a Poisson process with intensity $\lambda Y_{\alpha_1}(t_1)$. The conditional independence follows by similar arguments, and hence the first implication is proved.

Conversely, by [36], to prove Proposition 5.2 it is sufficient to prove (5.8). Denote by

$$\begin{aligned}\mathcal{H}^1_{s_1,s_2}&=\sigma(N_{\alpha_1,\alpha_2}(s_1+t,s)-N_{\alpha_1,\alpha_2}(s_1,s),t\geq 0,s\leq s_2)\\\mathcal{H}^2_{s_1,s_2}&=\sigma(N_{\alpha_1,\alpha_2}(s,s_2+t)-N_{\alpha_1,\alpha_2}(s,s_2),t\geq 0,s\leq s_1),\end{aligned}$$

so that $\mathcal{F}^{N_{\alpha_1,\alpha_2}}_{\infty,s_2}=\sigma(\mathcal{F}^{N_{\alpha_1,\alpha_2}}_{s_1,s_2},\mathcal{H}^1_{s_1,s_2})$ and $\mathcal{F}^{N_{\alpha_1,\alpha_2}}_{s_1,\infty}=\sigma(\mathcal{F}^{N_{\alpha_1,\alpha_2}}_{s_1,s_2},\mathcal{H}^2_{s_1,s_2})$. Then, denoting by $X \perp\!\!\!\perp Y|W$ the conditional independence of X and Y, given W, we have by hypothesis that

$$\mathcal{H}^1_{s_1,s_2}\perp\!\!\!\perp\mathcal{H}^2_{s_1,s_2}|\sigma(\mathcal{G},\mathcal{F}^{N_{\alpha_1,\alpha_2}}_{s_1,s_2}),\qquad\mathcal{H}^1_{s_1,s_2}\perp\!\!\!\perp\mathcal{F}^{N_{\alpha_1,\alpha_2}}_{s_1,s_2}|\mathcal{G},\qquad\mathcal{H}^2_{s_1,s_2}\perp\!\!\!\perp\mathcal{F}^{N_{\alpha_1,\alpha_2}}_{s_1,s_2}|\mathcal{G},$$

for any (s_1,s_2). Thus,

- $\mathcal{H}^2_{t_1,t_2}\perp\!\!\!\perp\mathcal{F}^{N_{\alpha_1,\alpha_2}}_{t_1,t_2},\mathcal{H}^1_{t_1,t_2}|\mathcal{G},$, $\mathcal{F}^{N_{\alpha_1,\alpha_2}}_{t_1,t_2}\perp\!\!\!\perp\mathcal{H}^1_{t_1,t_2}|\mathcal{G},$, $\mathcal{H}^1_{t_1,s_2}\subseteq\mathcal{H}^1_{t_1,t_2}$, $\mathcal{H}^2_{s_1,t_2}\subseteq\mathcal{H}^2_{t_1,t_2}$, then

  $$\mathrm{E}\Big(N_{\alpha_1,\alpha_2}(t_1,t_2)\Big|\sigma(\mathcal{F}^{N_{\alpha_1,\alpha_2}}_{\infty,s_2}\vee\mathcal{F}^{N_{\alpha_1,\alpha_2}}_{s_1,\infty},\mathcal{G})\Big)=\mathrm{E}\Big(N_{\alpha_1,\alpha_2}(t_1,t_2)\Big|\sigma(\mathcal{F}^{N_{\alpha_1,\alpha_2}}_{t_1,s_2}\vee\mathcal{F}^{N_{\alpha_1,\alpha_2}}_{s_1,t_2},\mathcal{G})\Big),$$

  and hence

  $$\mathrm{E}\Big(\Delta_{s_1,s_2}N_{\alpha_1,\alpha_2}(t_1,t_2)\Big|\sigma(\mathcal{F}^{N_{\alpha_1,\alpha_2}}_{\infty,s_2}\vee\mathcal{F}^{N_{\alpha_1,\alpha_2}}_{s_1,\infty},\mathcal{G})\Big)=\mathrm{E}\Big(\Delta_{s_1,s_2}N_{\alpha_1,\alpha_2}(t_1,t_2)\Big|\sigma(\mathcal{F}^{N_{\alpha_1,\alpha_2}}_{t_1,s_2}\vee\mathcal{F}^{N_{\alpha_1,\alpha_2}}_{s_1,t_2},\mathcal{G})\Big);\tag{5.10}$$

- note that $\mathcal{F}^{N_{\alpha_1,\alpha_2}}_{t_1,s_2}=\sigma(\mathcal{F}^{N_{\alpha_1,\alpha_2}}_{s_1,s_2},\mathcal{H})$, where $\mathcal{H}=\sigma(\Delta_{s_1,s_2}N_{\alpha_1,\alpha_2}(u,v),s_1\leq u\leq t_1,v\leq s_2)$. In addition, $\mathcal{H}^1_{s_1,t_2}\perp\!\!\!\perp\mathcal{F}^{N_{\alpha_1,\alpha_2}}_{s_1,t_2}|\mathcal{G}$, and $\sigma(\Delta_{s_1,s_2}N_{\alpha_1,\alpha_2}(t_1,t_2),\mathcal{H})\subseteq\mathcal{H}^1_{s_1,t_2}$. Hence

  $$\mathrm{E}\Big(\Delta_{s_1,s_2}N_{\alpha_1,\alpha_2}(t_1,t_2)\Big|\sigma(\mathcal{F}^{N_{\alpha_1,\alpha_2}}_{t_1,s_2}\vee\mathcal{F}^{N_{\alpha_1,\alpha_2}}_{s_1,t_2},\mathcal{G})\Big)=\mathrm{E}\Big(\Delta_{s_1,s_2}N_{\alpha_1,\alpha_2}(t_1,t_2)\Big|\sigma(\mathcal{H},\mathcal{G})\Big);\tag{5.11}$$

- now, note that both $N_{\alpha_1,\alpha_2}(t_1,t_2)-N_{\alpha_1,\alpha_2}(t_1,s_2)$ and $N_{\alpha_1,\alpha_2}(s_1,t_2)-N_{\alpha_1,\alpha_2}(s_1,s_2)$ belong to $\mathcal{H}^2_{t_1,s_2}$, while $\mathcal{H}\subseteq\mathcal{F}^{N_{\alpha_1,\alpha_2}}_{t_1,s_2}$. Hence

  $$\mathrm{E}\Big(\Delta_{s_1,s_2}N_{\alpha_1,\alpha_2}(t_1,t_2)\Big|\sigma(\mathcal{H},\mathcal{G})\Big)=\mathrm{E}(\Delta_{s_1,s_2}N_{\alpha_1,\alpha_2}(t_1,t_2)|\mathcal{G}).\tag{5.12}$$

Combining (5.10), (5.11) and (5.12) we finally get (5.8):

$$\begin{aligned}\mathrm{E}\Big(\Delta_{s_1,s_2}N_{\alpha_1,\alpha_2}(t_1,t_2)\Big|\sigma(\mathcal{F}^{N_{\alpha_1,\alpha_2}}_{\infty,s_2}\vee\mathcal{F}^{N_{\alpha_1,\alpha_2}}_{s_1,\infty},\mathcal{G})\Big)&=\mathrm{E}(\Delta_{s_1,s_2}N_{\alpha_1,\alpha_2}(t_1,t_2)|\mathcal{G})\\&=\lambda(Y_{\alpha_1}(t_1)-Y_{\alpha_1}(s_1))(Y_{\alpha_2}(t_2)-Y_{\alpha_2}(s_2)).\quad\square\end{aligned}$$

Theorem 5.2 shows an example of characterizations of FPRF. In [19], the authors proved a characterization of the inhomogeneous Poisson processes on the plane thorough its realizations on increasing families of points (called increasing path) and increasing families of sets, called increasing set (see also [2, 22]).

We are going to characterize an FPRF in the same spirit.

Let $\mathcal{A}$ be the collection of the closed rectangles $\{A_{t_1,t_2} : t \in \mathbb{R}^2_+\}$, where $A_{t_1,t_2} = \{(s_1, s_2) \in \mathbb{R}^2_+ : 0 \le s_i \le t_i, i = 1, 2\}$. The family $\mathcal{A}$ generates a topology of closed sets $\widetilde{\mathcal{A}}(u)$, which is closed under finite unions and arbitrary intersections, called a *lower set* family (see, e.g., [1, 23]). In other words, when a point (t_1, t_2) belongs to a set $A \in \widetilde{\mathcal{A}}(u)$, all the rectangle A_{t_1,t_2} is contained in A:

$$A \in \widetilde{\mathcal{A}}(u) \qquad \Longleftrightarrow \qquad A_{t_1,t_2} \subseteq A, \forall (t_1, t_2) \in A.$$

A function $\Gamma : \mathbb{R}_+ \to \widetilde{\mathcal{A}}(u)$ is called an *increasing set* if $\Gamma(0) = \{(0, 0)\}$, it is continuous, it is non-decreasing ($s \le t \Longrightarrow \Gamma(s) \subseteq \Gamma(t)$), and the area it underlies is finite for any t and goes to infinity when t increases ($\lim_{t\to+\infty} |\Gamma(t)| = \infty$). Note that, for a nonnegative measure μ on $B_{\mathbb{R}_+\times\mathbb{R}_+}$, it is well-defined the non-decreasing right-continuous function:

$$(\mu \circ \Gamma)(t) = \mu(\Gamma(t)).$$

Given an increasing path Γ and a random nonnegative measure N (in [23], it is an increasing and additive process), we may define the one-parameter process $N \circ \Gamma$ as the trace of N along Γ:

$$(N \circ \Gamma)(t) = N(\Gamma(t)), \quad t \ge 0.$$

THEOREM 5.3. *A random simple locally finite counting measure N_{α_1,α_2} is a FPRF iff, conditioned on $\mathcal{G}$, $N \circ \Gamma$ is an inhomogeneous Poisson process with intensity $\lambda(\mu_{\boldsymbol{\alpha}} \circ \Gamma)$, for any increasing set Γ, independent of $\mathcal{G}$.*

PROOF. Assume that N_{α_1,α_2} is a FPRF. Then, for any $0 \le s_1 < t_1 \le s_2 < t_2 \le \cdots \le s_n < t_n$, the sets $B_i = \Gamma(t_i) \setminus \Gamma(s_i)$ are disjoint. By (5.9),

$$\begin{aligned}
\mathrm{P}\Big(\bigcap_{i=1}^n \{(N \circ \Gamma)(s_i, t_i] = x_i\}\Big|\mathcal{G}\Big) &= \mathrm{P}\Big(\bigcap_{i=1}^n \{N_{\alpha_1,\alpha_2}(B_i) = x_i\}\Big|\mathcal{G}\Big) \\
&= \prod_{i=1}^n \frac{\exp(-\lambda\mu_{\boldsymbol{\alpha}}(B_i))(\lambda\mu_{\boldsymbol{\alpha}}(B_i))^{x_i}}{x_i!} \\
&= \prod_{i=1}^n \frac{\exp\big(-\lambda\cdot(\mu_{\boldsymbol{\alpha}} \circ \Gamma)(s_i, t_i]\big)\big(\lambda\cdot(\mu_{\boldsymbol{\alpha}} \circ \Gamma)(s_i, t_i]\big)^{x_i}}{x_i!}.
\end{aligned}$$

Conversely, note that that (5.9) may be checked only on disjoint rectangles $B_1, B_2, \ldots, B_n$ (see also [23]). After ordering partially the rectangles with respect to $\prec$, one can build an increasing sets Γ such that $B_i = \Gamma(t_i) \setminus \Gamma(s_i)$, where $0 \le s_1 < t_1 \le s_2 < t_2 \le \cdots \le s_n < t_n$. By hypothesis, $N \circ \Gamma$ is an inhomogeneous Poisson process with intensity $\mu_{\boldsymbol{\alpha}} \circ \Gamma$. Then,

$$\begin{aligned}
\mathrm{P}\Big(\bigcap_{i=1}^n \{N_{\alpha_1,\alpha_2}(B_i) = x_i\}\Big|\mathcal{G}\Big) &= \mathrm{P}\Big(\bigcap_{i=1}^n \{(N \circ \Gamma)(s_i, t_i] = x_i\}\Big|\mathcal{G}\Big) \\
&= \prod_{i=1}^n \frac{\exp\big(-\lambda\cdot(\mu_{\boldsymbol{\alpha}} \circ \Gamma)(s_i, t_i]\big)\big(\lambda\cdot(\mu_{\boldsymbol{\alpha}} \circ \Gamma)(s_i, t_i]\big)^{x_i}}{x_i!}
\end{aligned}$$

$$= \prod_{i=1}^{n} \frac{\exp(-\lambda\mu_{\boldsymbol{\alpha}}(B_i))(\lambda\mu_{\boldsymbol{\alpha}}(B_i))^{x_i}}{x_i!}. \square$$

Now, a function $\Gamma : \mathbb{R}_+ \to \mathbb{R}_+^2$ is called an *increasing path* if $\Gamma(0) = (0,0)$, it is continuous, it is non-decreasing ($s \le t \Longrightarrow \Gamma_1(s) \le \Gamma_1(t), \Gamma_2(s) \le \Gamma_2(t)$), and the area it underlies goes to infinity ($\lim_{t\to+\infty} \Gamma_1(t)\Gamma_2(t) = \infty$). In other words, an increasing path is an increasing set where, for each t, $\Gamma(t)$ is a rectangle. Given an increasing path Γ and a process $N(t_1, t_2)$, we define the one-parameter process $N \circ \Gamma$ as the trace of N along Γ:

$$(N \circ \Gamma)(t) = N(\Gamma_1(t), \Gamma_2(t)), \quad t \ge 0.$$

When dealing with the laws of the traces of a process along increasing paths, one cannot hope to prove, for instance, the conditional independence of two filtrations as ${\mathcal{H}^{\infty}}_{s_1,s_2}$ and ${\mathcal{H}^{\in}}_{s_1,s_2}$, since the event that belong to those filtrations are generated by the increments of the process on rwgions that are not comparable with respect to the partial order $\prec$.

As an example, there is no increasing path that separates the three rectangles $B_1 = \{(1,0) \prec z \prec (2,1)\}$, $B_2 = \{(0,1) \prec z \prec (1,2)\}$ and $B_3 = \{(1,1) \prec z \prec (2,2)\}$ and hence we cannot give the joint law of $\Delta_{(1,0)}N(2,1)$ and $\Delta_{(0,1)}N(1,2)$. On the other hand, Proposition 5.2 suggests that, if we assume the independence of $N(B_1)$ and $N(B_2)$ conditioned on $\mathcal{F}_{1,1}$, (5.9) may be proved for B_1, B_2 and B_3 via increasing paths (as in [2, 3, 19, 22]). This consideration has suggesed the following definition.

We say that the filtration satisfies the conditional independence condition or the Cairoli-Walsh condition ((F4) in [14], see also [25]) if for any $\mathcal{F}$-measurable integrable random variable Z, and for any (t_1, t_2) :

$$\mathrm{E}(\mathrm{E}(Z|\mathcal{F}_{t_1,\infty})|\mathcal{F}_{\infty,t_2}) = \mathrm{E}(\mathrm{E}(Z|\mathcal{F}_{\infty,t_2})|\mathcal{F}_{t_1,\infty}) = \mathrm{E}(Z|\mathcal{F}_{t_1,t_2}).$$

Thus, following the same ideas as in [2, 3, 19, 22], one can prove the following result.

THEOREM 5.4. *A random simple locally finite counting measure N_{α_1,α_2} is a FPRF iff, conditioned on $\mathcal{G}$, the Cairoli-Walsh condition holds and $N \circ \Gamma$ is an inhomogeneous Poisson process with intensity $Y_{\alpha_1}(\Gamma_1(t)) \cdot Y_{\alpha_2}(\Gamma_2(t))$, for any increasing path Γ.*

A remark on Set-Indexed Fractional Poisson Process. Let T be a metric space equipped with a Radon measure on its Borel sets. We assume existence of an indexing collection $\mathcal{A}$ on T, as it is defined in [23]. We are interested to considering processes indexed by a class of closed sets from T.

$\Gamma : \mathbb{R}_+ \to \mathcal{A}$ is called an increasing path if it is continuous and increasing: $s < t \Longrightarrow \Gamma(s) \subseteq \Gamma(t)$ (called a flow in [18])

We can now define Set-Indexed Fractional Poisson process.

A set-indexed process $X = \{X_U, U \in \mathcal{A}\}$ is called a Set-Indexed Fractional Poisson process(SIFPP), if for any increasing path Γ the process $X^{\Gamma} = \{X_{\Gamma(t)}, t \ge 0\}$ is an FPP.

Following results of [23], we can state

THEOREM 5.5. *Any SIFPP is a set-indexed Lévy process.*

Details and martingale characterizations will be presented elsewhere.

5.4. *Gergely-Yezhow characterization.* Let $(U_n, n \geq 1)$ be a sequence of i.i.d. $(0,1)$-uniform distributed random variables, independent of the processes Y_{α_i}, $i = 1, 2$. The random indexes associated to the 'records' $(\nu_n, n \geq 1)$ are inductively defined by

$$\nu_1(\omega) = 1, \qquad \nu_{n+1}(\omega) = \inf\{k > \nu_n(\omega) \colon U_k(\omega) > U_{\nu_n(\omega)}(\omega)\}.$$

It is well known that $P(\cap_n\{\nu_n < \infty\}) = 1$, and hence the k-th record V_k of the sequence is well defined: $V_0 := 0$, $V_k = U_{\nu_k}$. Since $V_n \geq \max(U_1, \dots, U_n)$, then $P(V_n \to 1) = 1$. Moreover, the number of U_n's that realize the maximum by time n is almost surely asymptotic to $\log(n)$ as $n \to \infty$. In other words, the sequence $(\nu_n)_n$ growths exponentially fast.

Now, given a increasing set Γ, we define

$$Y^{\Gamma}_t = \sum_n n \mathbb{1}_{[V_n, V_{n+1})}(1 - \exp(-\mu_{\boldsymbol{\alpha}} \circ \Gamma(t))) = \sup\{n \colon V_n \leq 1 - \exp(-\mu_{\boldsymbol{\alpha}} \circ \Gamma(t))\}.$$

THEOREM 5.6. *A random simple locally finite counting measure N_{α_1,α_2} is a FPRF iff $N \circ \Gamma$ is distributed as Y^{Γ}, for any increasing set Γ.*

PROOF. In the proof we assume that $\lim_t \mu_{\boldsymbol{\alpha}} \circ \Gamma(t) = \infty$ almost surely. When this is not the case, the proof should be changed as in [16], where generalized random variables are introduced exactly when $1 - \exp(-$"intensity at ∞"$) < 1$.

By Theorem 5.3, we must prove that, conditioned on $\mathcal{G}$, Y^{Γ} is an inhomogeneous Poisson process with intensity $\mu_{\boldsymbol{\alpha}} \circ \Gamma$. Conditioned on $\mathcal{G}$, let $F(t) := 1 - \exp(-\mu_{\boldsymbol{\alpha}} \circ \Gamma(t))$ be the continuous deterministic cumulative distribution function. Let F^- be its pseudo-inverse, and define $\xi_n = F^-(U_n)$, for each n. Then $(\xi_n, n \geq 1)$ is a sequence of i.i.d. random variables with cumulative function F. As in [16], put $\zeta'_n = \max(\xi_1, \dots, \xi_n)$, $(n = 1, 2, \dots)$ omitting in the increasing sequence

$$\zeta'_1, \zeta'_2, \dots, \zeta'_n, \dots$$

all the repeating elements except one, we come to the strictly increasing sequence [16, Eq. (3)]

$$\zeta_1, \zeta_2, \dots, \zeta_n, \dots$$

Now, since F^- is monotone, it is obvious by definition that $\zeta_n = F^-(V_n)$. Again, F^- is monotone, and hence

$$\begin{aligned} Y^{\Gamma}_t &= \sum_n n \mathbb{1}_{[F^-(V_n), F^-(V_{n+1}))}(F^-(1 - \exp(-\mu_{\boldsymbol{\alpha}} \circ \Gamma(t)))) \\ &= \sum_n n \mathbb{1}_{[\zeta_n, \zeta_{n+1})}(t), \end{aligned}$$

that is the process $v(t)$ defined in [16, Eq. (7')]. The thesis is now an application of [16, Theorem 1] and Theorem 5.3. □

5.5. *Random time change.* The process $\mu_{\boldsymbol{\alpha}}$ may be used to reparametrize the time of the increasing paths and sets. In fact, for any increasing path $\Gamma = (\Gamma_1(t), \Gamma_2(t))$, let

$$T(s,\omega) = \begin{cases} \inf\{t \colon Y_{\alpha_1}(\Gamma_1(t)) \cdot Y_{\alpha_2}(\Gamma_2(t))(\omega) > s\} & \text{if } \{t \colon Y_{\alpha_1}(\Gamma_1(t)) \cdot Y_{\alpha_2}(\Gamma_2(t))(\omega) > s\} \neq \varnothing; \\ \infty & \text{otherwise;} \end{cases}$$

be the first time that the intensity is seen to be bigger than s on the increasing path, and define

$$\Gamma_{\mu_{\boldsymbol{\alpha}}}(s,\omega) = \Gamma(T(s,\omega)) \tag{5.13}$$

the reparametrization of Γ made by $\mu_{\boldsymbol{\alpha}}$. Analogously, for any increasing set Γ, let

$$\Gamma_{\mu_{\boldsymbol{\alpha}}}(s,\omega) = \Gamma(\inf\{t\colon (\mu_{\boldsymbol{\alpha}}(\omega)\circ\Gamma)(t) > s\}).$$

We note that, for any fixed s and $A \in \widetilde{\mathcal{A}}(u)$

$$\{\omega\colon A \not\subseteq \Gamma_{\mu_{\boldsymbol{\alpha}}}(s)\} = \cup_{t\in\mathbb{Q}}\Big(\{A \not\subseteq \Gamma(t)\} \cap \{\mu_{\boldsymbol{\alpha}}(\Gamma(t)\cap A) \geq s\}\Big) \in \mathcal{G}_A, \tag{5.14}$$

where $\mathcal{G}_A = \sigma(\mu_{\boldsymbol{\alpha}}(A'), A' \subseteq A)$. We recall that a random measurable set $Z : \Omega \to \widetilde{\mathcal{A}}(u)$ is called a $\mathcal{G}_A$-stopping set if $\{A \subseteq Z\} \in \mathcal{G}_A$ for any A. As a consequence, the reparametrization given in (5.13) transforms $\Gamma(\cdot)$ into $\Gamma_{\mu_{\boldsymbol{\alpha}}}(\cdot)$, a family of continuous increasing stopping set by (5.14). Such a family is called an optional increasing set. The random time change theorem (which can be made an easy consequence of the characterization of the Poisson process given in [49]) together with Theorem 5.3 and Theorem 5.4 give the following corollaries, that can be seen as extensions of some results in [2, 3].

COROLLARY 5.2. *A random simple locally finite counting measure N_{α_1,α_2} is a FPRF iff, conditioned on $\mathcal{G}$, $N\circ\Gamma_{\mu_{\boldsymbol{\alpha}}}$ is a standard Poisson process, for any increasing set Γ.*

COROLLARY 5.3. *A random simple locally finite counting measure N_{α_1,α_2} is a FPRF iff, conditioned on $\mathcal{G}$, the Cairoli-Walsh condition holds [14, 25] and $N\circ\Gamma_{\mu_{\boldsymbol{\alpha}}}$ is a standard Poisson process, for any increasing path Γ.*

6. Fractional Differential Equations. A direct calculation may be applied to show that the marginal distribution of the classical Poisson random field $N(t_1,t_2),\ (t_1,t_2)\in\mathbb{R}^2_+$

$$p^c_k(t_1,t_2) = \mathrm{P}\left(N(t_1,t_2)=k\right) = \frac{e^{-\lambda t_1 t_2}(\lambda t_1 t_2)^k}{k!}, k = 0,1,2\ldots$$

satisfy the following differential-difference equations:

$$\frac{\partial^2 p^c_0(t_1,t_2)}{\partial t_1\,\partial t_2} = \left(-\lambda+\lambda^2 t_1 t_2\right) p^c_0(t_1,t_2); \tag{6.1}$$

$$\frac{\partial^2 p^c_1(t_1,t_2)}{\partial t_1\,\partial t_2} = \left(-3\lambda+\lambda^2 t_1 t_2\right) p^c_1(t_1,t_2) + \lambda p^c_0(t_1,t_2); \tag{6.2}$$

$$\frac{\partial^2 p^c_k(t_1,t_2)}{\partial t_1\,\partial t_2} = \left(-\lambda+\lambda^2 t_1 t_2\right) p^c_k(t_1,t_2) + \left(\lambda-2\lambda^2 t_1 t_2\right) p^c_{k-1}(t_1,t_2) + \lambda^2 p^c_{k-2}(t_1,t_2);\ k\geq 2; \tag{6.3}$$

and the initial conditions:

$$p^c_0(0,0)=1,\ p^c_k(0,0) = p^c_k(t_1,0) = p^c_k\ (0,t_2) = 0,\ k\geq 1.$$

We are now ready to derive the governing equations of the marginal distributions of FPRF $N_{\alpha_1,\alpha_2}(t_1,t_2),\ (t_1,t_2)\in\mathbb{R}^2_+$:

$$p^{\alpha_1,\alpha_2}_k(t_1,t_2) = \mathrm{P}\left(N_{\alpha_1,\alpha_2}(t_1,t_2)=k\right),\ k=0,1,2,\ldots \tag{6.4}$$

given by (5.2) or (5.3). These equations have something in common with the governing equations for the non-homogeneous Fractional Poisson processes [30].

For a function $u(t_1,t_2),\ (t_1,t_2)\in\mathbb{R}^2_+$, the Caputo-Djrbashian mixed fractional derivative of order $\alpha_1,\alpha_2\in(0,1)\times(0,1)$ is defined by

$$\mathrm{D}^{\alpha_1,\alpha_2}_{t_1,t_2}u(t_1,t_2)=\frac{1}{\Gamma(1-\alpha_1)\Gamma(1-\alpha_2)}\int_0^{t_1}\int_0^{t_2}\frac{\partial^2 u(\tau_1,\tau_2)}{\partial\tau_1\,\partial\tau_2}\frac{d\tau_1\,d\tau_2}{(t_1-\tau_1)^{\alpha_1}(t_2-\tau_2)^{\alpha_2}}$$
$$=\frac{1}{\Gamma(1-\alpha_1)\Gamma(1-\alpha_2)}\int_0^{t_1}\int_0^{t_2}\frac{\partial^2 u(t_1-\upsilon_1,t_2-\upsilon_2)}{\partial\upsilon_1\,\partial\upsilon_2}\frac{d\upsilon_1\,d\upsilon_2}{\upsilon_1^{\alpha_1}\upsilon_2^{\alpha_2}}.$$

Assuming that

$$e^{-s_1t_1-s_2t_2}\frac{\partial^2 u(t_1-\upsilon_1,t_2-\upsilon_2)}{\partial\upsilon_1\,\partial\upsilon_2}\upsilon_1^{-\alpha_1}\upsilon_2^{-\alpha_2}$$

is integrable as function of four variables $t_1,t_2,\upsilon_1,\upsilon_2$, the double Laplace transform

$$\mathcal{L}\left\{\mathrm{D}^{\alpha_1,\alpha_2}_{t_1,t_2}u(t_1,t_2);s_1,s_2\right\}=\int_0^\infty\int_0^\infty e^{-s_1t_1-s_2t_2}\mathrm{D}^{\alpha_1,\alpha_2}_{t_1,t_2}u(t_1,t_2)dt_1\,dt_2$$
$$=s_1^{\alpha_1}s_2^{\alpha_2}\tilde{u}(s_1,s_2)-s_1^{\alpha_1-1}s_2^{\alpha_2}\tilde{u}(s_1,0)-s_1^{\alpha_1}s_2^{\alpha_2-1}\tilde{u}(0,s_2)+s_1^{\alpha_1-1}s_2^{\alpha_2-1}\tilde{u}(0,0),\quad(6.5)$$

where $\tilde{u}(s_1,s_2)=\mathcal{L}\{u(t_1,t_2);s_1,s_2\}$ is the double Laplace transform of the function $u(t_1,t_2)$.

REMARK. Note that the Laplace transform of $f_\alpha(t,x)$ given by (2.4) as $\alpha=1$ is of the form e^{-sx} and its inverse is the delta distribution $\delta(t-x)$. Accordingly, as $\alpha\to1$, $f_\alpha(t,x)$ converges weakly to $\delta(t-x)$, and we denote it by $f_\alpha(t,x)\to\delta(t-x)$.

The proof of (6.5) is standard and we omit it (see [34, p. 37] for the one-dimensional case).

THEOREM 6.1. *Let $N(t_1,t_2)$, $(t_1,t_2)\in\mathbb{R}^2_+,\alpha_1,\alpha_2\in(0,1)\times(0,1)$, be the FPRF defined by (5.1).*

1) *Then its marginal distribution given in (6.4) satisfy the following fractional differential-integral recurrent equations:*

$$\mathrm{D}^{\alpha_1,\alpha_2}_{t_1,t_2}p_0^{\alpha_1,\alpha_2}(t_1,t_2)=\int_0^\infty\int_0^\infty\left(-\lambda+\lambda^2x_1x_2\right)p_0^{\alpha_1,\alpha_2}(x_1,x_2)\,f_{\alpha_1}(t_1,x_1)f_{\alpha_2}(t_2,x_2)dx_1dx_2;\quad(6.6)$$

$$\mathrm{D}^{\alpha_1,\alpha_2}_{t_1,t_2}p_1^{\alpha_1,\alpha_2}(t_1,t_2)=\int_0^\infty\int_0^\infty\Big[\left(-3\lambda+\lambda^2x_1x_2\right)p_1^{\alpha_1,\alpha_2}(x_1,x_2)$$
$$+\lambda p_0^{\alpha_1,\alpha_2}(x_1,x_2)\Big]f_{\alpha_1}(t_1,x_1)f_{\alpha_2}(t_2,x_2)dx_1dx_2;\quad(6.7)$$

$$\mathrm{D}^{\alpha_1,\alpha_2}_{t_1,t_2}p_k^{\alpha_1,\alpha_2}(t_1,t_2)=\int_0^\infty\int_0^\infty\Big[\left(-\lambda+\lambda^2x_1x_2\right)p_k^{\alpha_1,\alpha_2}(x_1,x_2)$$
$$+\left(\lambda-2\lambda^2x_1x_2\right)p_{k-1}^{\alpha_1,\alpha_2}(x_1,x_2)+\lambda^2x_1xp_{k-2}^{\alpha_1,\alpha_2}(x_1,x_2)\Big]$$
$$\times f_{\alpha_1}(t_1,x_1)f_{\alpha_2}(t_2,x_2)dx_1dx_2,\qquad k\geq2;\quad(6.8)$$

with the initial conditions:

$$p_0^{\alpha_1,\alpha_2}(0,0)=1,\ p_k^{\alpha_1,\alpha_2}(0,0)=p_k^{\alpha_1,\alpha_2}(t_1,0)=p_k^{\alpha_1,\alpha_2}(0,t_2)=0,\ k\geq1.$$

2) *For $\alpha_1\to1,\alpha_2\to1,f_{\alpha_1}(t_1,x_1)\to\delta(t_1-x_1)$, $f_{\alpha_2}(t_2,x_2)\to\delta(t_2-x_2)$, hence (6.6), (6.7) and (6.8) become (6.1), (6.2) and (6.3) correspondingly.*

PROOF. **1)** The initial conditions are easily checked using the fact that $Y_{\alpha_1}(0)=Y_{\alpha_2}(0)=0$ a.s. Let $p_k^{\alpha_1,\alpha_2}(t_1,t_2),k=0,1,2,\ldots$, be defined as in equations (5.2) or (5.3). Then the characteristic function of the FPRF, for $z\in\mathbb{R}$:

$$\hat{p}(t_1,t_2;z)=\mathrm{E}\exp\{izN_{\alpha_1,\alpha_2}(t_1,t_2)\}=\int_0^\infty\int_0^\infty e^{\lambda x_1x_2(e^{iz}-1)}f_{\alpha_1}(t_1,x_1)f_{\alpha_2}(t_2,x_2)dx_1dx_2.\quad(6.9)$$

Taking the double Laplace transform of (6.9) and using (2.4) and (5.4) yields

$$\bar{p}(s_1,s_2;z)=\widetilde{\hat{p}}(t_1,t_2;z)=\int_0^\infty\int_0^\infty e^{-s_1t_1-s_2t_2}\hat{p}(t_1,t_2;z)dt_1dt_2 \tag{6.10}$$
$$=s_1^{\alpha_1-1}s_2^{\alpha_2-1}\int_0^\infty\int_0^\infty e^{\lambda x_1x_2(e^{iz}-1)}e^{s_1^{\alpha_1-1}s_2^{\alpha_2-1}}f_{\alpha_1}(t_1,x_1)f_{\alpha_2}(t_2,x_2)dx_1dx_2,$$

and

$$\bar{p}\,(0,0,z)=\bar{p}\,(0,s_2,z)=\bar{p}\,(s_1,0,z)=0.$$

Using an integration by parts for a double integral [28]:

$$\int_0^\infty\int_0^\infty F(x_1,x_2)H\,(dx_1,dx_2)=\int_0^\infty\int_0^\infty H\,([x_1,\infty)\times[x_2,\infty))\,F\,(dx_1,dx_2)$$
$$+\int_0^\infty H\,([x_1,\infty)\times[0,\infty))\,F\,(dx_1,0)$$
$$+\int_0^\infty H\,([0,\infty)\times[x_2,\infty))\,F\,(0,dx_2)+F(0,0)H\,([0,\infty)\times[0,\infty))\,,$$

we get from (6.5), (6.10) and (6.10) with

$$F(x_1,x_2)=\exp\left\{\lambda x_1x_2(e^{iz}-1)\right\},\ H\,(dx_1,dx_2)=\exp\left\{-s_1^{\alpha_1}x_1-s_2^{\alpha_2}x_2\right\}dx_1dx_2,$$
$$\bar{p}(s_1,s_2;z)=s_1^{\alpha_1-1}s_2^{\alpha_2-1}\Big[\int_0^\infty\int_0^\infty\frac{\partial^2\exp\left\{ix_1x_2(e^{iz}-1)\right\}}{\partial x_1\,\partial x_2}$$
$$\times\frac{\exp\left\{-s_1^{\alpha_1}x_1-s_2^{\alpha_2}x_2\right\}}{s_1^{\alpha_1}s_2^{\alpha_2}}dx_1,dx_2+\frac{\hat{p}(0,0,z)}{s_1^{\alpha_1}s_2^{\alpha_2}}\Big].$$

Thus

$$s_1^{\alpha_1}s_2^{\alpha_2}\bar{p}(s_1,s_2;z)-\hat{p}(0,0,z)$$
$$=s_1^{\alpha_1-1}s_2^{\alpha_2-1}\int_0^\infty\int_0^\infty\frac{\partial^2\exp\left\{ix_1x_2(e^{iz}-1)\right\}}{\partial x_1\,\partial x_2}\exp\left\{-s_1^{\alpha_1}x_1-s_2^{\alpha_2}x_2\right\}dx_1,dx_2$$

Using (6.5), (2.4) we can invert the double Laplace transform as follows:

$$\mathrm{D}^{\alpha_1,\alpha_2}_{t_1,t_2}\hat{p}\,(t_1,t_2,z)=\int_0^\infty\int_0^\infty\frac{\partial^2\exp\left\{ix_1x_2(e^{iz}-1)\right\}}{\partial x_1\,\partial x_2}f_{\alpha_1}(t_1,x_1)f_{\alpha_2}(t_2,x_2)dx_1dx_2.$$

And finally, by inverting the characteristic function (6.9), we obtain

$$\mathrm{D}^{\alpha_1,\alpha_2}_{t_1,t_2}\hat{p}\,(t_1,t_2,z)\,p_k^{\alpha_1,\alpha_2}\,(t_1,t_2)=\int_0^\infty\int_0^\infty\left[\frac{\partial^2}{\partial x_1\,\partial x_2}p_k^c(x_1,x_2)\right]f_{\alpha_1}(t_1,x_1)f_{\alpha_2}(t_2,x_2)dx_1dx_2.$$

Using (6.1), (6.2) and (6.3) we arrive to (6.6), (6.7) and (6.8) correspondingly.

2) Finally, as $\alpha_j\to1, j=1,2$ we have $e^{-s_j^{\alpha_j}x_j}\to e^{-s_jx_j}, j=1,2$, and their Laplace inversions are delta function: $\delta(t_j-x_j), j=1,2$. Thus, 2) is proven. □



(a) $\alpha_1 = 0.95$, $\alpha_2 = 0.5$, $\lambda = 100$



(b) $\alpha_1 = \alpha_2 = 0.75$, $\lambda = 100$



(c) $\alpha_1 = 0.9$, $\alpha_2 = 0.75$, $\lambda = 100$



(d) $\alpha_1 = \alpha_2 \sim 1$, $\lambda = 10000$

Fig 1: Simulations of the inverse stable subordinators $Y^{(1)}_{\alpha_1}(t)$ and $Y^{(2)}_{\alpha_2}(t)$ and the corresponding FPRF N_{α_1,α_2} for different values of α_1 and α_2. Top-left: simulation of $Y^{(1)}_{\alpha_1}(t)$, top-right: simulation of $Y^{(2)}_{\alpha_2}(t)$, bottom-(left-right): simulation of N_{α_1,α_2}, the rotation shows the connection with marginal intensity

7. Simulations. In this section we show some simulations of FPRF made with Matlab based on the α-stable random number generator function `stblrnd`.

The subordinators L_α are simulated exactly at times $t_n = n\Delta$, where $\Delta = 0.0005$ till they reach a defined value S_{end}. More precisely,

$$L_\alpha(0) = 0; \qquad L_\alpha(t_n) = L_\alpha(t_{n-1}) + X, \qquad n = 1, 2, \dots, N$$

where X is independently simulated with `stblrnd`$(\alpha,\ 1,\ \sqrt[\alpha]{\Delta}\ ,\ 0)$. Accordingly,

$$\mathrm{E}e^{-sX} = \exp\{-(s\sqrt[\alpha]{\Delta})^\alpha\} = \exp\{-\Delta s^\alpha\}, \qquad s \geq 0,$$

and hence

$$\mathrm{E}e^{-sL_\alpha(t_n)} = \exp\{-t_n s^\alpha\}, \qquad s \geq 0, n = 0, 1, \dots, N.$$

The simulation of the inverse stable subordinators $Y_\alpha(s), s \in [0, T_{\text{end}}]$ are thus made at times $s_n = L_\alpha(t_n), n = 1, \dots, N$ with values $Y_\alpha(s_n) = n\Delta$.

To simulate a FPRF $N_{\alpha_1,\alpha_2}(s^1, s^2)$ on the window $(0, S_{\text{end}}) \times (0, S_{\text{end}})$, we first simulate two independent inverse stable subordinators $Y^{(1)}_{\alpha_1}(s^1_n), n = 1, \dots, N_1$ and $Y^{(2)}_{\alpha_2}(s^2_n), n = 1, \dots, N_2$.

By Proposition 5.2, the value of N_{α_1,α_2} on each rectangle $(s^1_n, s^1_{n+1}) \times (s^1_n, s^1_{n+1})$ is a Poisson random variable with mean Δ^2. As $\Delta^2 \ll 1$, we approximate it with a Bernoulli random variable Y of parameter Δ^2. When $Y = 1$, we add a point at random inside the rectangle.

In Figure 1 the simulations of the inverse stable subordinators $Y^{(1)}_{\alpha_1}(t)$ and $Y^{(2)}_{\alpha_2}(t)$ and the corresponding FPRF N_{α_1,α_2} for different values of α_1 and α_2 are shown. The simulations of N_{α_1,α_2} are plotted twice: we have rotated each figure in order to underline the spatial dependence of the spread of the points of the process N_{α_1,α_2} in connection with the marginal intensities $Y^{(1)}_{\alpha_1}(t)$ and $Y^{(2)}_{\alpha_2}(t)$. For example, in Figure 1(c) two different marginal distribution are expected since $\alpha_1 = 0.9$ and $\alpha_2 = 0.75$. While $Y^{(1)}_{0.9}(t)$ produces a quite uniform distribution of points, $Y^{(2)}_{0.75}(t)$ generates clusters in correspondence of its steeper slopes.

With a MonteCarlo procedure, we also compute

$$p_k(t_1, t_2) = \mathrm{P}\left(N(Y_1(t_1), Y_2(t_2)) = k\right)$$

for different values of t_1, t_2, α_1 and α_2. In fact, in (5.2), we approximate $f_{\alpha_1}(t_1, x_1) f_{\alpha_2}(t_2, x_2) dx_1 dx_2$ with

$$g(x_1, x_2) = \frac{1}{N^2} \sum_{n_1=1}^{N} \sum_{n_2=1}^{N} \mathbb{1}_{X_{n_1}}(x_1) \mathbb{1}_{Y_{n_2}}(x_2)$$

where $(X_n, n = 1, \dots, N)$ and $(Y_n, n = 1, \dots, N)$ are independent sequences of i.i.d. distributed as $Y^{(1)}_{\alpha_1}(t_1)$ and $Y^{(2)}_{\alpha_2}(t_2)$, respectively. The simulations with $N = 1500$ are presented in Figure 2. We underline the variety of the shape of distributions that can be generated with this two-parameter model in addition to its flexibility to include, for example, different cluster phenomena.

Acknowledgements. N. Leonenko and E. Merzbach wish to thank G. Aletti for two visits to University of Milan.

N. Leonenko was supported in particular by Cardiff Incoming Visiting Fellowship Scheme and International Collaboration Seedcorn Fund and Australian Research Council's Discovery Projects funding scheme (project number DP160101366).

The authors thank anonymous reviewers, for their helpful comments.



(a) $t_1 = t_2 = 5$, $\alpha_1 = 0.5$, $\alpha_2 = .75$

(b) $t_1 = t_2 = 5$, $\alpha_1 = 0.95$, $\alpha_2 = .075$



(c) $t_1 = t_2 = 5$, $\alpha_1 = \alpha_2 = 0.75$



(d) $t_1 = 3$, $t_2 = 7$, $\alpha_1 = \alpha_2 = 0.75$

Fig 2: Simulations of the distribution of $Y^{(1)}_{\alpha_1}(t_1)$, $Y^{(2)}_{\alpha_2}(t_2)$ and the corresponding $p_k(t_1, t_2) = \mathrm{P}\left(N(Y_1(t_1), Y_2(t_2)) = k\right)$ for $\lambda = 10$ and different values of t_1, t_2, α_1 and α_2.

APPENDIX A: COVARIANCE STRUCTURE OF PARAMETER-CHANGED POISSON RANDOM FIELDS

In this Appendix, we prove a general result that can be used to compute the covariance structure of the parameter-changed Poisson random field:

$$Z(t_1,t_2) = N(Y_1(t_1), Y_2(t_2)),\ (t_1,t_2) \in \mathbb{R}^2_+,$$

where $Y_1 = \{Y_1(t_1), t_1 \geq 0\}$ and $Y_2 = \{Y_2(t_2), t_2 \geq 0\}$ are independent non-negative non-decreasing stochastic processes, in general non-Markovian with non-stationary and non-independent increments, and $N = \{N(t_1,t_2), (t_1,t_2) \in \mathbb{R}^2_+\}$ is a PRF with intensity $\lambda > 0$. We also assume that Y_1 and Y_2 are independent of N.

For example, Y_1 and Y_2 might be inverse subordinators.

THEOREM A.1. *Suppose that N is a PRF, Y_1 and Y_2 are two non-decreasing non-negative independent stochastic processes which are also independent of N. Then*

1) if $\mathrm{E}Y_1(t_1) = U_1(t_1)$ and $\mathrm{E}Y_2(t_2) = U_2(t_2)$ exist, then $\mathrm{E}Z(t_1,t_2)$ exists and

$$\mathrm{E}Z(t_1,t_2) = \mathrm{E}N(1,1)\mathrm{E}Y_1(t_1)\mathrm{E}Y_2(t_2);$$

2) if Y_1 and Y_2 have second moments, so does Z and

$$\begin{aligned}\mathrm{Var}Z(t_1,t_2) = [\mathrm{E}N(1,1)]^2 \left\{\mathrm{E}Y_1^2(t_1)\mathrm{E}Y_2^2(t_2) - \left(\mathrm{E}Y_1^2(t_1)\right)^2 \left(\mathrm{E}Y_2^2(t_2)\right)^2\right\}\\ +\mathrm{Var}N(1,1)\mathrm{E}Y_1(t_1)\mathrm{E}Y_2(t_2)\end{aligned}$$

and its covariance function

$$\mathrm{Cov}(Z(t_1,t_2), Z(s_1,s_2)) = \mathrm{Cov}\left(N(Y_1(t_1), Y_2(t_2)), N(Y_1(s_1), Y_2(s_2))\right) \tag{A.1}$$

for $s_1 < t_1, s_2 < t_2$ is given by:

$$\begin{aligned}&[\mathrm{E}N(1,1)]^2 \{\mathrm{Cov}\left(Y_1(t_1), Y_1(s_1)\right)\mathrm{Cov}\left(Y_2(t_2), Y_2(s_2)\right) +\\ &\quad + \left(\mathrm{E}Y_2(t_2)\mathrm{E}Y_2(s_2) - \mathrm{E}Y_2^2(s_2)\right)\left(\mathrm{Cov}\left(Y_1(t_1), Y_1(s_1)\right) - \mathrm{E}Y_1^2(s_1)\right)\\ &\quad + \left(\mathrm{E}Y_1(t_1)\mathrm{E}Y_1(s_1) - \mathrm{E}Y_1^2(s_1)\right)\left(\mathrm{Cov}\left(Y_2(t_2), Y_2(s_2)\right) - \mathrm{E}Y_2^2(s_2)\right)\}\\ &\quad + \mathrm{Var}N(1,1)\mathrm{E}Y_1(s_1)\mathrm{E}Y_2(s_2)\end{aligned}$$

and for any (s_1,s_2), and (t_1,t_2) from $\mathbb{R}^2_+$

$$\begin{aligned}&[\mathrm{E}N(1,1)]^2 \Big\{\mathrm{Cov}\left(Y_1(t_1), Y_1(s_1)\right)\mathrm{Cov}\left(Y_2(t_2), Y_2(s_2)\right)\\ &\quad + \left(\mathrm{E}Y_2(t_2)\mathrm{E}Y_2(s_2) - \mathrm{E}Y_2^2(\min(s_2,t_2))\right)\left(\mathrm{Cov}\left(Y_1(t_1), Y_1(s_1)\right) - \mathrm{E}Y_1^2(\min(s_1,t_1))\right)\\ &\quad + \left(\mathrm{E}Y_1(t_1)\mathrm{E}Y_1(s_1) - \mathrm{E}Y_1^2(\min(s_1,t_1))\right)\left(\mathrm{Cov}\left(Y_2(t_2), Y_2(s_2)\right) - \mathrm{E}Y_2^2(\min(s_2,t_2))\right)\Big\}\\ &\quad + \mathrm{Var}N(1,1)\mathrm{E}Y_1(\min(s_1,t_1))\mathrm{E}Y_2(\min(s_2,t_2)).\end{aligned}$$

REMARK. These formulae are valid for any Lévy random field $N = \{N(t_1,t_2),(t_1,t_2) \in \mathbb{R}^2_+\}$, with finite expectation $\mathrm{E}N(1,1)$ and finite variance $\mathrm{Var}N(1,1)$, for PRF $\mathrm{E}N(1,1) = \lambda$; $\mathrm{Var}N(1,1) = \lambda$ and to apply these formulae one needs to know

$$U_1(t_1) = \mathrm{E}Y_1(t),\ U_2(t_2) = \mathrm{E}Y_2(t),\ U_1^{(2)}(t_1) = \mathrm{E}Y_1^2(t),\ U_2^{(2)}(t_1) = \mathrm{E}Y_2^2(t),$$

and $\mathrm{Cov}\left(Y_1(t_1), Y_1(s_1)\right)$, $\mathrm{Cov}\left(Y_2(t_2), Y_2(s_2)\right)$ which are available for many non-negative processes $Y_1(t)$ and $Y_2(t)$ induction inverse subordinators.

REMARK. One can compute the following expression for the one-dimensional distribution of the parameter-changed PRF:

$$\begin{aligned}\mathrm{P}\left(N(Y_1(t_1),Y_2(t_2))=k\right) &= p_k(t_1,t_2)\\ &= \int_0^\infty\int_0^\infty \frac{e^{-\lambda x_1x_2}(\lambda x_1x_2)^k}{k!} f_1(t_1,x_1)f_2(t_2,x_2)dx_1dx_2,\ k=0,1,2,\dots\end{aligned}$$

where

$$f_i(t_i,x_i)=\frac{d}{dx_i}\mathrm{P}\left\{Y_i(t_i)\le x_i\right\}=\frac{d}{dx_i}G^{(i)}_{t_i}(x_i),\qquad i=1,2.$$

and its Laplace transform:

$$\mathcal{L}\left\{p_k(t_1,t_2);s_1,s_2\right\}=\int_0^\infty\int_0^\infty \frac{e^{-\lambda x_1x_2}(\lambda x_1x_2)^k}{k!}\mathcal{L}\left\{f_1(t_1,x_1);s_1\right\}\mathcal{L}\left\{f_2(t_2,x_2);s_2\right\}dx_1dx_2,$$

where

$$\mathcal{L}\left\{f_i(t_i,x_i);s_i\right\}=\int_0^\infty e^{-s_it_i}f_i(t_i,x_i)dt_i,\qquad i=1,2.$$

PROOF OF THEOREM A.1. We denote

$$\begin{aligned}&G^{(1)}_{t_1}(u_1)=\mathrm{P}\left\{Y_1(t_1)\le u_1\right\},\qquad &&G^{(1)}_{t_1,s_1}(u_1,v_1)=\mathrm{P}\left\{Y_1(t_1)\le u_1,Y_1(s_1)\le v_1\right\},\\ &G^{(2)}_{t_2}(u_2)=\mathrm{P}\left\{Y_2(t_2)\le u_2\right\},\qquad &&G^{(2)}_{t_2,s_2}(u_2,v_2)=\mathrm{P}\left\{Y_2(t_2)\le u_2,Y_2(s_2)\le v_2\right\}.\end{aligned}$$

We know that for a PRF

$$\mathrm{E}\Delta_{s_1,s_2}N(t_1,t_2)=\left(\mathrm{E}N(1,1)\right)^2(t_1-s_1)(t_2-s_2)=\mathrm{Var}\Delta_{s_1,s_2}N(t_1,t_2);$$

$$\mathrm{E}\left[\Delta_{s_1,s_2}N(t_1,t_2)\right]^2=\mathrm{E}N(1,1)(t_1-s_1)(t_2-s_2)+\left[\mathrm{E}N(1,1)(t_1-s_1)(t_2-s_2)\right]^2.$$

To prove 1) we use simple conditioning arguments:

$$\mathrm{E}Z(t_1,t_2)=\int_0^\infty\int_0^\infty u\ v\ \mathrm{E}N(1,1)G^{(1)}_{t_1}(du)G^{(2)}_{t_2}(dv)=\mathrm{E}N(1,1)\mathrm{E}Y_1(t_1)\mathrm{E}Y_2(t_2).$$

Let us prove 2).
For the variance, we have

$$\begin{aligned}\mathrm{Var}Z(t_1,t_2) &= \mathrm{E}\left(Z(Y_1(t_1),Y_2(t_2)\right)^2-\left(\mathrm{E}Z(Y_1(t_1),Y_2(t_2)\right)^2\\ &= \int_0^\infty\int_0^\infty\left(\mathrm{E}N^2(u_1,u_2)+\mathrm{Var}N(u_1,u_2)\right)G^{(1)}_{t_1}(du_1)G^{(2)}_{t_2}(du_2)\\ &\quad -\left(\mathrm{E}N(1,1)\mathrm{E}Y_1(t_1)\mathrm{E}Y_2(t_2)\right)^2\\ &= \int_0^\infty\int_0^\infty\left[\left(\mathrm{E}N(1,1)\right)^2u_1^2,u_2^2+\mathrm{Var}N(1,1)u_1,u_2\right]G^{(1)}_{t_1}(du_1)G^{(2)}_{t_2}(du_2)\\ &\quad -\left(\mathrm{E}N(1,1)\mathrm{E}Y_1(t_1)\mathrm{E}Y_2(t_2)\right)^2\\ &= \left(\mathrm{E}N(1,1)\right)^2\mathrm{E}Y_1^2(t_1)\mathrm{E}Y_2^2(t_2)+\mathrm{Var}N(1,1)\mathrm{E}Y_1(t_1)\mathrm{E}Y_2(t_2)\\ &\quad -\left(\mathrm{E}N(1,1)\mathrm{E}Y_1(t_1)\mathrm{E}Y_2(t_2)\right)^2\\ &= \left(\mathrm{E}N(1,1)\right)^2\left\{\mathrm{E}Y_1^2(t_1)\mathrm{E}Y_2^2(t_2)-\left(\mathrm{E}Y_1(t_1)\right)^2\left(\mathrm{E}Y_2(t_2)\right)^2\right\}\\ &\quad -\mathrm{Var}N(1,1)\mathrm{E}Y_1(t_1)\mathrm{E}Y_2(t_2).\end{aligned}$$

To compute the covariance structure, first we consider the case when $s_1 < t_1,\ s_2 < t_2$: Then

$$
\begin{aligned}
&\mathrm{E}N(s_1,s_2)N(t_1,t_2)\\
&= \mathrm{E}\{N(t_1,t_2) - N(t_1,s_2) - N(t_2,s_1) + N(s_1,s_2) + N(t_1,s_2) + N(t_2,s_1)\\
&\quad - N(s_1,s_2)\}N(s_1,s_2)\\
&= \mathrm{E}\Delta_{s_1,s_2}N(t_1,t_2)\mathrm{E}N(s_1,s_2) + \mathrm{E}N(t_1,s_2)N(s_1,s_2) + \mathrm{E}N(t_2,s_1)N(s_1,s_2) - \mathrm{E}N^2(s_1,s_2)\\
&= (t_1-s_1)(t_2-s_2)\left[\mathrm{E}N(1,1)\right]^2 s_1s_2 + \mathrm{E}\left\{N(t_1,s_2) - N(t_1,s_1) + N(s_1,s_2)\right\}N(s_1,s_2)\\
&\quad + \mathrm{E}\left\{N(t_2,s_1) - N(s_1,s_2) + N(s_1,s_2)\right\}N(s_1,s_2) - \mathrm{E}N^2(s_1,s_2)\\
&= \left[\mathrm{E}N(1,1)\right]^2 (t_1-s_1)(t_2-s_2)s_1s_2 + \mathrm{E}N^2(s_1,s_2) + \mathrm{E}N^2(s_1,s_2) - \mathrm{E}N^2(s_1,s_2)\\
&= \left[\mathrm{E}N(1,1)\right]^2 (t_1-s_1)(t_2-s_2)s_1s_2 + \mathrm{Var}N^2(s_1,s_2) + \left[\mathrm{E}N(s_1,s_2)\right]^2\\
&= \left[\mathrm{E}N(1,1)\right]^2 (t_1-s_1)(t_2-s_2)s_1s_2 + \mathrm{Var}N(1,1)s_1s_2 + \left[\mathrm{E}N(1,1)\right]^2 s_1^2s_2^2\\
&= \left[\mathrm{E}N(1,1)\right]^2 \left[(t_1-s_1)(t_2-s_2)s_1s_2 + s_1^2s_2^2\right] + s_1^2s_2^2\mathrm{Var}N(1,1).
\end{aligned}
$$

Note that

$$
(t_1-s_1)(t_2-s_2)s_1s_2 + s_1^2s_2^2 = t_1t_2s_1s_2 - t_1s_1s_2^2 - t_2s_1^2s_2 + 2s_1^2s_2^2.
$$

Since the processes N, Y_1, Y_2 are independent, a conditioning argument yields

$$
\begin{aligned}
&\mathrm{E}N(Y_1(t_1),Y_2(t_2))N(Y_1(s_1),Y_2(s_2))\\
&\qquad = \left[\mathrm{E}N(1,1)\right]^2 \Bigg\{\int_0^\infty\int_0^\infty\int_0^\infty\int_0^\infty u_1u_2v_1v_2G^{(1)}_{t_1,s_1}(du_1,dv_1)G^{(2)}_{t_2,s_2}(du_2,dv_2)\\
&\qquad\quad - \int_0^\infty\int_0^\infty\int_0^\infty\int_0^\infty u_1v_1v_2^2G^{(1)}_{t_1,s_1}(du_1,dv_1)G^{(2)}_{t_2}(du_2)G^{(2)}_{s_2}(dv_2)\\
&\qquad\quad - \int_0^\infty\int_0^\infty\int_0^\infty\int_0^\infty u_2v_1^2v_2G^{(1)}_{t_2,s_2}(du_2,dv_2)G^{(1)}_{t_1,}(du_1)G^{(1)}_{s_1}(dv_1)\\
&\qquad\quad + 2\int_0^\infty\int_0^\infty\int_0^\infty\int_0^{2\infty} v_1^2v_2^2G^{(1)}_{t_1,}(du_1)G^{(1)}_{s_1}(dv_1)G^{(2)}_{t_2}(du_2)G^{(2)}_{s_2}(dv_2)\Bigg\}\\
&\qquad\quad + \mathrm{Var}N(1,1)\int_0^\infty\int_0^\infty\int_0^\infty\int_0^\infty v_1v_2G^{(1)}_{t_1}(du_1)G^{(1)}_{s_1}(dv_1)G^{(2)}_{t_2}(du_2)G^{(2)}_{s_2}(dv_2)\\
&\qquad = \left[\mathrm{E}N(1,1)\right]^2 \left\{\mathrm{E}Y_1(t_1)Y_1(s_1)\mathrm{E}Y_2(t_2)Y_2(s_2) - \mathrm{E}Y_1(t_1)Y_1(s_1)\mathrm{E}Y_2^2(s_2)\right.\\
&\qquad\quad \left. - \mathrm{E}Y_2(t_2)Y_2(s_2)\mathrm{E}Y_2^2(s_1) + 2\mathrm{E}Y_2^2(s_1)Y_2^2(s_2)\right\} + \mathrm{Var}N(1,1)\mathrm{E}Y_1(s_1)\mathrm{E}Y_2(s_2),
\end{aligned}
$$

and

$$
\begin{aligned}
&\mathrm{Cov}\left[N(Y_1(t_1),Y_2(t_2)),N(Y_1(s_1),Y_2(s_2))\right]\\
&= \mathrm{E}N(Y_1(t_1),Y_2(t_2))N(Y_1(s_1),Y_2(s_2)) - \left[\mathrm{E}N(1,1)\right]^2\mathrm{E}Y_1(t_1)\mathrm{E}Y_2(t_2)\mathrm{E}Y_1(s_1)\mathrm{E}Y_2(s_2)\\
&\quad = \left[\mathrm{E}N(1,1)\right]^2\left\{\mathrm{E}Y_1(t_1)Y_1(s_1)\mathrm{E}Y_2(t_2)Y_2(s_2) - \mathrm{E}Y_1(t_1)Y_1(s_1)\mathrm{E}Y_2^2(s_2) - \mathrm{E}Y_2(t_2)Y_2(s_2)\mathrm{E}Y_2^2(s_1)\right.\\
&\quad \left. + 2\mathrm{E}Y_1^2(s_1)\mathrm{E}Y_2^2(s_2) - \mathrm{E}Y_1(t_1)\mathrm{E}Y_1(s_1)\mathrm{E}Y_2(t_2)\mathrm{E}Y_2(s_2)\right\} + \mathrm{Var}N(1,1)\mathrm{E}Y_1(s_1)\mathrm{E}Y_2(s_2)\\
&= \left[\mathrm{E}N(1,1)\right]^2\left\{\left[\mathrm{Cov}(Y_1(t_1),Y_1(s_1)) + \mathrm{E}Y_1(t_1)\mathrm{E}Y_1(s_1)\right]\left[\mathrm{Cov}(Y_2(t_2),Y_2(s_2)) + \mathrm{E}Y_2(t_2)\mathrm{E}Y_2(s_2)\right]\right.\\
&\quad - \left[\mathrm{Cov}(Y_1(t_1),Y_1(s_1)) + \mathrm{E}Y_1(t_1)\mathrm{E}Y_1(s_1)\right]\mathrm{E}Y_2^2(s_2) - \left[\mathrm{Cov}(Y_2(t_2),Y_2(s_2))\right.\\
&\quad \left.\left. + \mathrm{E}Y_2(t_2)\mathrm{E}Y_2(s_2)\right]\mathrm{E}Y_1^2(s_1)
\end{aligned}
$$

$$
\begin{aligned}
&+ 2\mathrm{E}Y_1^2(s_1)\mathrm{E}Y_2^2(s_2) - \mathrm{E}Y_1(t_1)\mathrm{E}Y_1(s_1)\mathrm{E}Y_2(t_2)\mathrm{E}Y_2(s_2)\big\} + \mathrm{Var}N(1,1)\mathrm{E}Y_1(s_1)\mathrm{E}Y_2(s_2)\\
&= [\mathrm{E}N(1,1)]^2\,\{\mathrm{Cov}(Y_1(t_1),Y_1(s_1))\mathrm{Cov}(Y_2(t_2),Y_2(s_2)) + \mathrm{Cov}(Y_1(t_1),Y_1(s_1))\mathrm{E}Y_2(t_2)\mathrm{E}Y_2(s_2)\\
&\quad + \mathrm{Cov}(Y_2(t_2),Y_2(s_2))\mathrm{E}Y_1(t_1)\mathrm{E}Y_1(s_1)\\
&\quad - \mathrm{Cov}(Y_1(t_1),Y_1(s_1))\mathrm{E}Y_2^2(s_2) - \mathrm{E}Y_1(t_1)\mathrm{E}Y_1(s_1)\mathrm{E}Y_2^2(s_2)\\
&\quad - \mathrm{Cov}(Y_2(t_2),Y_2(s_2))\mathrm{E}Y_1^2(s_1) - \mathrm{E}Y_2(t_2)\mathrm{E}Y_2(s_2)\mathrm{E}Y_1^2(s_1)\\
&\quad + 2\mathrm{E}Y_1^2(s_1)\mathrm{E}Y_2^2(s_2) - \mathrm{E}Y_1(t_1)\mathrm{E}Y_1(s_1)\mathrm{E}Y_2(t_2)\mathrm{E}Y_2(s_2)\big\} + \mathrm{Var}N(1,1)\mathrm{E}Y_1(s_1)\mathrm{E}Y_2(s_2)\\
&= [\mathrm{E}N(1,1)]^2\,\{\mathrm{Cov}(Y_1(t_1),Y_1(s_1))\mathrm{Cov}(Y_2(t_2),Y_2(s_2)) + \mathrm{Cov}(Y_1(t_1),Y_1(s_1))[\mathrm{E}Y_2(t_2)\mathrm{E}Y_2(s_2)\\
&\quad - \mathrm{E}Y_2^2(s_2)] + \mathrm{Cov}(Y_2(t_2),Y_2(s_2))\left[\mathrm{E}Y_1(t_1)\mathrm{E}Y_2(s_2) - \mathrm{E}Y_1^2(s_1)\right]\\
&\quad + \mathrm{E}Y_1^2(s)(\mathrm{E}Y_2^2(s_2) - \mathrm{E}Y_2(t_2)\mathrm{E}Y_2(s_2))\\
&\quad + \mathrm{E}Y_2^2(s_2)(\mathrm{E}Y_1^2(s_1) - \mathrm{E}Y_1(t_1)\mathrm{E}Y_1(s_1))\big\} + \mathrm{Var}N(1,1)\mathrm{E}Y_1(s_1)\mathrm{E}Y_2(s_2)\\
&= [\mathrm{E}N(1,1)]^2\,\big\{\mathrm{Cov}(Y_1(t_1),Y_1(s_1))\mathrm{Cov}(Y_2(t_2),Y_2(s_2))\left[\mathrm{E}Y_2(t_2)\mathrm{E}Y_2(s_2) - \mathrm{E}Y_2^2(s_2)\right]\\
&\quad \times \left(\mathrm{Cov}(Y_1(t_1),Y_1(s_1)) - \mathrm{E}Y_1^2(s_1)\right)\\
&\quad + \left(\mathrm{E}Y_1(t_1)\mathrm{E}Y_1(s_1)) - \mathrm{E}Y_1^2(s_1)\right)\left(\mathrm{Cov}(Y_2(t_2),Y_2(s_2)) - \mathrm{E}Y_2^2(s_2)\right)\big\}\\
&\quad + \mathrm{Var}N(1,1)\mathrm{E}Y_1(s_1)\mathrm{E}Y_2(s_2).
\end{aligned}
$$

In a similar way, one can consider the case $s_1 > t_1,\ s_2 < t_2$, where

$$
\begin{aligned}
\mathrm{E}N(t_1,t_2)N(s_1,s_2) &= -\mathrm{E}\left(N(s_1,t_2) - N(t_1,t_2) - N(s_1,s_2) + N(t_1,s_2)\right.\\
&\quad \left. -N(s_1,t_2) - N(t_1,s_2) + N(s_1,s_2)\right)N(s_1,s_2)\\
&= -\mathrm{E}\Delta_{t_1,s_2}N(s_1,t_2)N(s_1,s_2) + \mathrm{E}N(s_1,t_2)N(s_1,s_2)\\
&\quad -\mathrm{E}N^2(s_1,s_2) + \mathrm{E}N(t_1,s_2)N(s_1,s_2)
\end{aligned}
$$

and after routine calculations one can arrive at (A.1). □

Proof of Proposition 5.1. It follows from Theorem A.1 and Proposition 2.1. □

REFERENCES

[1] Aletti, G. (2001). On different topologies for set-indexing collections. *Statist. Probab. Lett.* **54** 67–73. MR1857872 (2002f:60064)

[2] Aletti, G. and Capasso, V. (1999). Characterization of spatial Poisson along optional increasing paths—a problem of dimension's reduction. *Statist. Probab. Lett.* **43** 343–347. MR1707943 (2000d:60084)

[3] Aletti, G. and Capasso, V. (2002). Reduction of dimension for spatial point processes and right continuous martingales. Characterization of spatial Poisson processes. *Stoch. Stoch. Rep.* **72** 1–9. MR1896435 (2003e:60101)

[4] Baddeley, A. (2007). Spatial point processes and their applications. In *Stochastic geometry. Lecture Notes in Math.* **1892** 1–75. Springer, Berlin. MR2327290 (2008c:60045)

[5] Beghin, L. (2012). Random-time processes governed by differential equations of fractional distributed order. *Chaos Solitons Fractals* **45** 1314–1327. MR2990245

[6] Beghin, L. and Orsingher, E. (2009). Fractional Poisson processes and related planar random motions. *Electron. J. Probab.* **14** 1790–1827. MR2535014 (2010m:60168)

[7] Beghin, L. and Orsingher, E. (2010). Poisson-type processes governed by fractional and higher-order recursive differential equations. *Electron. J. Probab.* **15** 684–709. MR2650778 (2011f:60168)

[8] Biermé, H., Demichel, Y. and Estrade, A. (2013). Fractional Poisson field and fractional Brownian field: why are they resembling but different? *Electron. Commun. Probab.* **18** 13. MR3033594

[9] Bingham, N. H. (1971). Limit theorems for occupation times of Markov processes. *Z. Wahrscheinlichkeitstheorie und Verw. Gebiete* **17** 1–22. MR0281255 (43 ##6974)

[10] Bowsher, C. G. and Swain, P. S. (2012). Identifying sources of variation and the flow of information in biochemical networks. *PNAS* **109** E1320-E1328.

[11] Brémaud, P. (1975). An extension of Watanabe's theorem of characterization of Poisson processes over the positive real half line. *J. Appl. Probability* **12** 396–399. MR0380970 (52 ##1867)

[12] Brémaud, P. (1981). *Point Processes and Queues*. Springer-Verlag, New York-Berlin. MR636252 (82m:60058)

[13] Busani, O. (2016). Aging uncoupled continuous time random walk limits. *Electron. J. Probab.* **21** paper no. 7, 17 pp.

[14] Cairoli, R. and Walsh, J. B. (1975). Stochastic integrals in the plane. *Acta Math.* **134** 111–183. MR0420845 (54 ##8857)

[15] Daley, D. J. (1999). The Hurst index of long-range dependent renewal processes. *Ann. Probab.* **27** 2035–2041. MR1742900 (2000k:60175)

[16] Gergely, T. and Yezhow, I. I. (1973). On a construction of ordinary Poisson processes and their modelling. *Z. Wahrscheinlichkeitstheorie und Verw. Gebiete* **27** 215–232. MR0359012 (50 ##11467)

[17] Haubold, H. J., Mathai, A. M. and Saxena, R. K. (2011). Mittag-Leffler functions and their applications. *J. Appl. Math.* Art. ID 298628, 51. MR2800586 (2012e:33061)

[18] Herbin, E. and Merzbach, E. (2013). The set-indexed Lévy process: stationarity, Markov and sample paths properties. *Stochastic Process. Appl.* **123** 1638–1670. MR3027894

[19] Ivanoff, B. G. and Merzbach, E. (1990). Characterization of compensators for point processes on the plane. *Stochastics Stochastics Rep.* **29** 395–405.

[20] Ivanoff, B. G. and Merzbach, E. (1994). A martingale characterization of the set-indexed Poisson process. *Stochastics Stochastics Rep.* **51** 69–82. MR1380763 (97c:60125)

[21] Ivanoff, B. G. and Merzbach, E. (2006). What is a multi-parameter renewal process? *Stochastics* **78** 411–441. MR2281679 (2008h:60354)

[22] Ivanoff, B. G., Merzbach, E. and Plante, M. (2007). A compensator characterization of point processes on topological lattices. *Electron. J. Probab.* **12** 47–74. MR2280258 (2008h:60187)

[23] Ivanoff, G. and Merzbach, E. (2000). *Set-Indexed Martingales. Monographs on Statistics and Applied Probability* **85**. Chapman & Hall/CRC, Boca Raton, FL. MR1733295 (2001g:60105)

[24] Kallenberg, O. (1997). *Foundations of modern probability*. Springer-Verlag, New York [etc.].

[25] Krengel, U. and Sucheston, L. (1981). Stopping rules and tactics for processes indexed by a directed set. *J. Multivariate Anal.* **11** 199–229. MR618785

[26] Leonenko, N. and Merzbach, E. (2015). Fractional Poisson fields. *Methodol. Comput. Appl. Probab.* **17** 155–168. MR3306677

[27] Leonenko, N. N., Meerschaert, M. M., Schilling, R. L. and Sikorskii, A. (2014). Correlation structure of time-changed Lévy processes. *Commun. Appl. Ind. Math.* **6** e-483, 22 pp. MR3277310

[28] Leonenko, N. N., Meerschaert, M. M. and Sikorskii, A. (2013). Fractional Pearson diffusions. *J. Math. Anal. Appl.* **403** 532–546. MR3037487

[29] Leonenko, N. N., Ruiz-Medina, M. D. and Taqqu, M. S. (2011). Fractional elliptic, hyperbolic and parabolic random fields. *Electron. J. Probab.* **16** 1134–1172. MR2820073 (2012m:60112)

[30] Leonenko, N. N., Scalas, E. and Trinh, M.-T. (2016). The fractional non-homogeneous Poisson process arXiv:1601.03965.

[31] Mainardi, F., Gorenflo, R. and Scalas, E. (2004). A fractional generalization of the Poisson processes. *Vietnam J. Math.* **32** 53–64. MR2120631

[32] Mainardi, F., Gorenflo, R. and Vivoli, A. (2005). Renewal processes of Mittag-Leffler and Wright type. *Fract. Calc. Appl. Anal.* **8** 7–38. MR2179226

[33] Meerschaert, M. M., Nane, E. and Vellaisamy, P. (2011). The fractional Poisson process and the inverse stable subordinator. *Electron. J. Probab.* **16** 1600–1620. MR2835248 (2012k:60252)

[34] Meerschaert, M. M. and Scheffler, H.-P. (2008). Triangular array limits for continuous time random walks. *Stochastic Process. Appl.* **118** 1606–1633. MR2442372 (2010b:60135)

[35] Meerschaert, M. M. and Sikorskii, A. (2012). *Stochastic Models for Fractional Calculus. de Gruyter Studies in Mathematics* **43**. Walter de Gruyter & Co., Berlin. MR2884383

[36] Merzbach, E. and Nualart, D. (1986). A characterization of the spatial Poisson process and changing time. *Ann. Probab.* **14** 1380–1390. MR866358 (88d:60145)

[37] Merzbach, E. and Shaki, Y. Y. (2008). Characterizations of multiparameter Cox and Poisson processes by the renewal property. *Statist. Probab. Lett.* **78** 637–642. MR2409527 (2009d:60151)

[38] Molchanov, I. and Ralchenko, K. (2015). A generalisation of the fractional Brownian field based on non-Euclidean norms. *J. Math. Anal. Appl.* **430** 262–278. MR3347213

[39] Molchanov, I. and Ralchenko, K. (2015). Multifractional Poisson process, multistable subordinator and related limit theorems. *Statist. Probab. Lett.* **96** 95–101. MR3281752

[40] Podlubny, I. (1999). *Fractional Differential Equations. Mathematics in Science and Engineering* **198**. Academic Press, Inc., San Diego, CA. MR1658022 (99m:26009)

[41] Repin, O. N. and Saichev, A. I. (2000). Fractional Poisson law. *Radiophys. and Quantum Electronics* **43** 738–741. MR1910034

[42] Ross, S. (2011). *A first course in probability*, Eight ed. Macmillan Co., New York; Collier Macmillan Ltd., London.

[43] Stoyan, D., Kendall, W. S. and Mecke, J. (1987). *Stochastic Geometry and its Applications. Wiley Series in Probability and Mathematical Statistics: Applied Probability and Statistics*. John Wiley & Sons, Ltd., Chichester With a foreword by D. G. Kendall. MR895588 (88j:60034a)

[44] Uchaikin, V. V. and Zolotarev, V. M. (1999). *Chance and stability. Modern Probability and Statistics*. VSP, Utrecht Stable distributions and their applications, With a foreword by V. Yu. Korolev and Zolotarev. MR1745764

[45] Veillette, M. and Taqqu, M. S. (2010). Numerical computation of first passage times of increasing Lévy processes. *Methodol. Comput. Appl. Probab.* **12** 695–729. MR2726540 (2012f:60150)

[46] Veillette, M. and Taqqu, M. S. (2010). Using differential equations to obtain joint moments of first-passage times of increasing Lévy processes. *Statist. Probab. Lett.* **80** 697–705. MR2595149 (2011c:60129)

[47] Wang, X.-T. and Wen, Z.-X. (2003). Poisson fractional processes. *Chaos Solitons Fractals* **18** 169–177. MR1984556 (2005b:60092)

[48] Wang, X.-T., Wen, Z.-X. and Zhang, S.-Y. (2006). Fractional Poisson process. II. *Chaos Solitons Fractals* **28** 143–147. MR2174587

[49] Watanabe, S. (1964). On discontinuous additive functionals and Lévy measures of a Markov process. *Japan. J. Math.* **34** 53–70. MR0185675 (32 ##3137)

E-mail: giacomo.aletti@unimi.it

E-mail: LeonenkoN@Cardiff.ac.uk

E-mail: ely.merzbach@biu.ac.il